

*Aquaculture pur la Santé*

# Native Prawns: A Candidate New Tool for Schistosomiasis Control

LABORATORY, FIELD, AND MODELING STUDIES

Giulio De Leo, & Susanne Sokolow,

ICTP – Sao Paulo

January 14, 2015

Email: [ssokolow@stanford.edu](mailto:ssokolow@stanford.edu), [deleo@stanford.edu](mailto:deleo@stanford.edu)



## Key partners

- **Sanna Sokolow** (Stanford & UCSB)
- **Chelsea Wood** (University of Michigan):
- **Kevin Lafferty, David Lopez-Carr, Armand Kuris** (UC Santa Barbara)
- **Shawn Coyle, Thomas Delomas, James Tidwell**, (Kentucky State University)
- **James Sanchirico** (UC Davis)
- EPLS field team (in addition to **Gilles Riveau** and **Nicolas**)
- *Members of the Gates/GCC Grants Advisory Board*
  - **Mike Hsieh** (Biomedical Research Institute)
  - **Artemis Koukounari** (King's College, London, UK)
  - **Sam Loker** (University of New Mexico)
  - **William Secor & Sue Montgomery** (CDC, Atlanta)
  - **David Rollinson** (London Natural History Museum, UK)
  - **Amir Sagi, Amit** (Ben-Gurion University of the Negev, Israel)
- **External collaborators:**
  - **Marino Gatto, Renato Casagrandi, Lorenzo Mari** (Politecnico di Milano, Italy)
  - **Andrea Rinaldo, Enrico Bertuzzo, Javier** (EPLS, Switzerland)
  - **Daniel Colley** (SCORE director, University of Georgia - external advisor)
  - Agence Nationale de l'Aquaculture (ANA) & Agency of Integration and Agricultural Development (Senegal)



WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission



**Sanna Sokolow**, PhD in Disease Ecology  
 - Co-PI with Armand Kuris (UCSB) of the NIH-  
 K08 project "Emergence and Biological  
 Control of Schistosomiasis" NIH Grant # PHS  
 K08AI082284, 2009-2012




## Senegal River Basin



WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission


## Diamina Dam, Senegal



WORKING PAPER - NOT FOR CITATION



The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission







WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission



WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission




WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission



## Schistosoma lifecycle



## Pathology

- Chronic anemia
- Growth stunting
- Wasting
- Cognitive impairment
- Lack of physical fitness
- Genital lesions
  - Increased risk of HIV
- Infertility
- Liver failure
- Bladder cancer



WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission

## Treatment options and eradication strategies



WORKING PAPER - NOT FOR CITATION


The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission





- **A Success: USAID's Neglected Tropical Disease Program**

“nearly 16 million people [affected by Schistosomiasis] were treated [with praziquantel] during the first year...55 million people by the third year”

“major drug donors...totaled more than \$1.4 billion...”






WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission

## Transmission sites




WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission




## Reinfection patterns of schistosomiasis after treatment

Webster et al 2013 *Acta tropica*





WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission



WORKING PAPER - NOT FOR CITATION


The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission



WORKING PAPER - NOT FOR CITATION


The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission





WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission



## LABORATORY STUDIES

Sokolow et al. *Acta Tropica* 2013

### Lab studies

*Summary of the main results*

**Previous studies demonstrated that *Macrobrachium rosenbergii* are predators of *B. glabrata***

- Roberts and Kuris 1990, Lee et al 1983
- Jordan, 1985 The St. Lucia Project

**Here, we show that the African native prawns *M. vollehovenii* :**

- **Are voracious snail predators**
- **Consume both *Biomphalaria* and *Bulinus* spp. snails**
- **Consume both infected and uninfected snails**
- **Exhibit snail consumption rates of 6 to 20 or more snails/prawn/day**

*depending on prawn and snail size*




WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission

- All the experiments shown in the following were aimed at understanding accurately **how many snails prawns eat per unit time** as a function of:
  - prawn size
  - snail size
  - density
  - species
  - infectious status
- Without these data, it is difficult to make accurate predictions

### Does predation efficiency depend upon prawn size?

Sokolow et al Acta Tropica 2013




WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission




## Predation depends on prawn size



Sokolow et al Acta Tropica 2013

## Do *small* prawns (<2-3 g) have any effect on *snail* population growth?

- Small prawns (<2 g) were set in tanks with large, pop. founder, egg-laying snails
- Then the emergence of offspring (“recruitment”) of:
  - small...
  - then medium
  - and finally large snails
 was monitored through time




Sokolow et al Acta Tropica 2013


WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission



## Small prawns can control snail population growth



How many average-size snails can  
prawns of different size eat?



- **What prawn size is most efficient in removing snail?**
- **Does the goal of harvesting large prawns conflict with the public health goal of eating as many snails as possible?**
- **If you can stock 1kg of prawns and want to control snail abundance, what is better?**  
to stock **lots of tiny prawns**  
or **few large ones?**




WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission



Are infected snails less likely to be predated than susceptible snails?

Prawns consumed more infected snails over uninfected



## Summary of Predation Results

- Native prawns are voracious predators of both snail species
- Consume both infected and uninfected snails
- Prawn consumption rate is predictable, i.e. btw 6-20 depending upon snail/prawn body size
- Fishery and public health goals are compatible


- So, prawns do eat snails in the lab experiments, but...
- Is there any evidence that they are doing it also in the wild?
- Does this have any measurable effect of disease transmission, in terms of prevalence and intensity of infection?



**FIELD TRIALS  
SENEGAL RIVER  
BASIN  
2011-2013**

```


graph TD
    A[Laboratory experiments  
What is the predation efficiency  
of native prawns on  
target snails?] --> B[Analysis of field data  
What is the evidence of  
predation in the wild?  
What is the impact on  
disease transmission?]
    B --> C[Scaling-up  
Modelling analysis  
Under what  
conditions prawns  
can contribute to  
disease control?]
    C --> D[What is the most  
effective control  
strategy?]
  
```



**Senegal River Delta**

**We present data from Lampsar 1 and 2, as it is a matched pair of villages that was monitored for the entire study period 2011-2013:**

- once **\*before\***
- three times **\*after\*** prawns were installed




## The net enclosure at Lampsar



WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission







What about human data?

Let's review the sampling scheme...

WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission



WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission

## Summary of Field Research Findings:

### 1. Dramatically Fewer Infectious/Shedding Snails at Prawn site

- Prawns can sustainably lower the density of infected & SHEDDING snails


### 2. Lower Human Re-infection Rate at Prawn Site

- Prawns protect against re-infection after PZQ treatment

### 3. Lower Urine Egg Burdens For Those At Prawn Site

- Prawns reduce egg burdens, after PZQ treatment

- Why only few shedding snails at the prawns site?
- Still high reinfection rates at the prawn site: how many prawns do we need to stock to have a better control of the disease?
- If both MDA and prawns stocking are used is the effect additive or synergetic?



## Mathematical models of Schistosomiasis

McDonald 1965

Woolhouse 1991 *Acta tropica*

$$\frac{dm}{dt} = \alpha Ny - \gamma m$$

$$\frac{dy}{dt} = \beta Hm(1 - y) - \mu'' y$$

Mean  
number of  
parasites  
per person

prevalence  
of infected  
snails

***m*: mean parasite burden**

***y*: disease prevalence in snail**

***N*: snail number, constant**

***H*: human population**


**No latent period in snails**

**No predation with functional response**

WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission






## RESULTS

WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission




Why the effect on abundance of shedding snails is stronger than the effect on total snail abundance?

- Schistosomiasis prepatent time is long and comparable to snail life expectancy
- Prawns reduces life expectancy of infective (*but not yet shedding snails*) just enough *to prevent them from shedding*
- Schisto reproductive number drops below 1 even though snail density remain significant
- If this is true, we shall expect a reduction not just in snail abundance but also in snail body size at the prawn site.

WORKING PAPER - NOT FOR CITATION


The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission

## Seasonality



Sturrock et al. 2001 Parasitology

## Simulations with seasonality in snail dynamics




WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission

## Summary of Model Results:

- Our model predictions fit well with observed patterns
- Restoring prawn population (whether through stocking or fish ladders) is likely to alleviate disease burden
- Stronger effect on:
  - # of shedding snails (vs. total snail pop.)
  - human egg burden (vs. prevalence of infection)


## Scenario analysis




**If both MDA and prawns stocking are used,  
is their effect additive or synergetic?**



## PZQ (80% coverage) vs “Prawns + PZQ”



## Treated vs. **untreated** people



WORKING PAPER - NOT FOR CITATION

The present report is not for citation, quotation or circulation as whole or in part without author's explicit permission

## Conclusions

- The combination of **prawn stocking** with **PZQ administration** can deliver results that no such a measure can achieve alone
- local elimination becomes a more likely outcome if control of the natural snail reservoir through restoration of the prawn pop. is integrated with PZQ administration and other control strategies (sanitation, outreach and education, etc. )

## Current and future developments

- Historical analysis of control
- World-wide geographical analysis of prawns, dams and schisto
- Gates– scale up
- CNH – socio-economic models/assessment (optimization)
- EEID (Jason Rohr) – agriculture and schisto
- EPFL (Andrea Rinaldo) – hydrology and schisto
- Sanna and I – more detailed simulations/ optimal control/etc.