



Higher Spins in Hyperspace

Dmitri Sorokin

INFN, Padova Section

Work in progress with Mirian Tsulaia (Canberra, Australia)

Sao Paulo, November 5, 2013

Motivation

- Different formulations of a theory are useful for revealing its different properties and features
 - Metric-like formulation
 - Frame-like formulation - Unfolded HS dynamics – Non-linear HS equations

4d HS fields:
$$\omega(x, y, \bar{y}) = \sum_{k,j=0}^{\infty} dx^m \omega_m^{\alpha_1 \dots \alpha_k \dot{\beta}_1 \dots \dot{\beta}_j}(x) y_{\alpha_1} \dots y_{\alpha_k} \bar{y}_{\dot{\beta}_1} \dots \bar{y}_{\dot{\beta}_j}$$

extension of 4d space-time with spinorial (twistor-like) directions

We are interested in a different hyperspace extension which also incorporates all 4d HS fields (an alternative to Kaluza-Klein). Free HS theory is a simple theory of a “hyper” scalar and spinor.

- Study of (hidden) symmetries can provide deeper insights into the structure of the theory and may help to find its most appropriated description
 - HS symmetries of HS theory are infinite-dimensional. They control in a very restrictive way the form of the unfolded equations and, hence, HS field interactions
- **Question:** what is the largest **finite-dimensional** symmetry of a HS system and can we learn something new from it?

Sp(8) symmetry of 4d HS theory (*Fronsdal, 1985*)

$$SO(1,3) \subset SO(2,3) \subset SO(2,4) \subset Sp(8)$$

$$SO(2,3) \approx Sp(4)$$

Sp(8) acts on infinite spectrum of 4d HS single-particle states ($s=0, 1/2, 1, 3/2, 2, \dots$)

this is a consequence of Flato-Fronsdal Theorem, 1978:

Tensor product of two 3d singleton moduli comprises all the massless fields of spin s in 4d

Singletons are 3d massless scalar and spinor fields which enjoy 3d conformal symmetry

$$SO(2,3) \approx Sp(4)$$

$$S \otimes S \Rightarrow Sp(4) \times Sp(4) \subset Sp(8)$$

Can Sp(8) play a role similar to Poincaré or conformal symmetry acting geometrically on a hyperspace containing 4d space-time?

Whether a 4d HS theory can be formulated as a field theory on this hyperspace?

`Geometrically' means: $\delta_{conf} x^m = a^m + l^m_n x^n + b x^m + k^m x^2 - 2k_n x^n x^m$

Fronsdal '85: minimal dimension of the Sp(8) hyperspace (containing 4d space-time) is 10

Particles and fields in Sp(8) hyperspace

- It took about 15 years to realize Fronsdal's idea in concrete terms

Bandos & Lukierski '98: Twistor-like (super) particle on a tensorial space
(their motivation was not related to HS theory, but to supersymmetry)

Most general $N=1$ susy in flat 4d: $\{Q_\alpha, Q_\beta\} = \gamma_{\alpha\beta}^m P_m + \gamma_{\alpha\beta}^{mn} Z_{mn}, \quad [P_m, Z_{nl}] = 0$

$$\gamma_{\alpha\beta}^m = \gamma_{\beta\alpha}^m, \quad \alpha, \beta = 1, 2, 3, 4$$

10d **space** $P_m \rightarrow x^m$ (4d coordinates), $Z_{mn} \rightarrow y^{mn} = -y^{nm}$ (6 extra coordinates)
coordinates: $X^{\alpha\beta} = X^{\beta\alpha} = \frac{1}{2} x^m \gamma_m^{\alpha\beta} + \frac{1}{4} y^{mn} \gamma_{mn}^{\alpha\beta}$ - 4×4 matrix coordinates

Superparticle action: $S = \int d\tau \lambda_\alpha \lambda_\beta (\dot{X}^{\alpha\beta} - i\theta^\alpha \dot{\theta}^\beta), \quad \lambda_\alpha$ - commuting twistor-like variable

possesses hidden (generalized superconformal) symmetry $Osp(1|8) \supset Sp(8)$

Quantization (*Bandos, Lukierski & D.S. '99*)

$$\left(\frac{\partial}{\partial X^{\alpha\beta}} - i\lambda_\alpha \lambda_\beta \right) \Phi(X, \lambda) = 0$$

- describes in 4d free fields of any spin $s=0, 1/2, 1, 3/2, 2, \dots$

Field theory in flat Sp(8) hyperspace

Field equations in flat hyperspace (*Vasiliev '01*):

Fourier transform $C(X, \xi) = \int d^4 \lambda e^{i \lambda_\alpha \xi^\alpha} \Phi(X, \lambda) \Rightarrow \left(\frac{\partial}{\partial X^{\alpha\beta}} + i \frac{\partial^2}{\partial \xi^\alpha \partial \xi^\beta} \right) C(X, \xi) = 0$ **Free unfolded equations**

$$C(X, \xi) = b(X) + \xi^\alpha f_\alpha(X) + \sum \xi^{\alpha_1} \dots \xi^{\alpha_k} C_{\alpha_1 \dots \alpha_k}(X)$$

$b(X)$ and $f_\alpha(X)$ are independent scalar and spinor hyperfields satisfying the equations:

$$\begin{aligned} (\partial_{\alpha\beta} \partial_{\gamma\delta} - \partial_{\alpha\gamma} \partial_{\beta\delta}) b(X) &= 0 \\ \partial_{\alpha\beta} f_\gamma(X) - \partial_{\alpha\gamma} f_\beta(X) &= 0 \end{aligned}$$

4d content of $b(X)$ and $f_\alpha(X)$ are Bargman-Wigner HS curvatures (*Vasiliev '01, Bandos et. al. '05*):

$$(X^{\alpha\beta} = X^{\beta\alpha} = \frac{1}{2} x^m \gamma_m^{\alpha\beta} + \frac{1}{4} y^{mn} \gamma_{mn}^{\alpha\beta})$$

Integer spins: $b(x^m, y^{mn}) = \varphi(x) + F_{mn}(x) y^{mn} + (R_{mn,pq}(x) - \frac{1}{2} \eta_{mp} \partial_n \partial_q \varphi) y^{mn} y^{pq} + \dots$

1/2 integer spins: $f^\alpha(x^m, y^{mn}) = \psi^\alpha(x) + (\Psi_{mn}^\alpha(x) - \frac{1}{2} \partial_m (\gamma_n \psi)^\alpha) y^{mn} + \dots$

Eoms and Bianchi: $\partial^2 \varphi = 0; \quad \partial_{[l} F_{mn]} = 0, \quad \partial^m F_{mn} = 0; \quad R_{[mn,p]q} = 0, \quad \eta^{mp} R_{mn,pq} = 0; \quad \dots$

Sp(8) transformations in hyperspace (symmetries of the field equations)

Conformal transformations: $\delta_{conf} x^m = a^m + l^m_n x^n + b x^m + k^m x^2 - 2k_n x^n x^m$

Sp(8) transformations: $\delta_{Sp(8)} X^{\alpha\beta} = a^{\alpha\beta} + 2g_{\gamma}^{(\alpha} X^{\beta)\gamma} - X^{\alpha\gamma} k_{\gamma\delta} X^{\delta\beta}$

$$\delta b = -\delta X^{\alpha\beta} \partial_{\alpha\beta} b - \frac{1}{2} (g_{\alpha}^{\alpha} - k_{\alpha\beta} X^{\alpha\beta}) b,$$

$$\delta f_{\gamma} = -\delta X^{\alpha\beta} \partial_{\alpha\beta} f_{\gamma} - \frac{1}{2} (g_{\alpha}^{\alpha} - k_{\alpha\beta} X^{\alpha\beta}) f_{\gamma} - (g_{\gamma}^{\alpha} - k_{\gamma\beta} X^{\beta\alpha}) f_{\alpha}$$

conformal weights of the fields

Sp(8) generators:

$$P_{\alpha\beta} = \frac{\partial}{\partial X^{\alpha\beta}}, \quad L_{\alpha}^{\beta} = 2X^{\beta\gamma} \frac{\partial}{\partial X^{\gamma\alpha}}, \quad K^{\alpha\beta} = X^{\alpha\gamma} X^{\beta\delta} \frac{\partial}{\partial X^{\gamma\delta}}$$

generators of GL(4)

$$[P, P] = 0, \quad [K, K] = 0, \quad [P, K] = L,$$

$$[L, L] = L, \quad [L, P] = P, \quad [L, K] = K$$

Hyperspace is a coset space: $P = \frac{Sp(8)}{GL(4) \times K}$

Hyperspace extension of $AdS(4)$

(Bandos, Lukierski, Preitschopf, D.S. '99; Vasiliev '01)

- 10d group manifold $Sp(4) \sim SO(2,3)$

$$AdS_4 = \frac{Sp(4)}{SO(1,3)} \quad Sp(4) = \frac{Sp(8)}{GL(4) \times K} = P + K \quad \text{- different } Sp(8) \text{ coset}$$

Like Minkowski and $AdS(4)$ spaces, which are conformally flat, the flat hyperspace and $Sp(4)$ are (locally) related to each other by a “generalized conformal” transformation

$Sp(2M)$ group manifolds are GL-flat (Plyushchay, D.S. & Tsulaia '03)

Algebra of covariant derivatives on $Sp(4)$:

$$[\nabla_{\alpha\beta}, \nabla_{\gamma\delta}] = \frac{1}{2r} (C_{\alpha(\gamma} \nabla_{\delta)\beta} + C_{\beta(\gamma} \nabla_{\delta)\alpha}), \quad C_{\alpha\beta} = -C_{\beta\alpha}, \quad r - \text{ is } Sp(4) \text{ (or AdS4) radius}$$

$$\nabla_{\alpha\beta} = G_{\alpha}^{\gamma} G_{\beta}^{\delta} \frac{\partial}{\partial X^{\gamma\delta}}, \quad G_{\alpha}^{\gamma}(X) = \delta_{\alpha}^{\gamma} + \frac{1}{4r} X_{\alpha}^{\gamma}; \quad \Omega^{\alpha\beta}(X) = G_{\gamma}^{-1\alpha} G_{\delta}^{-1\beta} dX^{\gamma\delta} \quad Sp(4) \text{ Cartan form}$$

GL-flatness is important for the relation between the field equations in flat and $Sp(4)$ hyperspace

HS field equations in Sp(4)

(Didenko and Vasiliev '03; Plyushchay, D.S. & Tsulaia '03)

Flat hyperspace equations:

$$(\partial_{\alpha\beta}\partial_{\gamma\delta} - \partial_{\alpha\gamma}\partial_{\beta\delta}) b(X) = 0$$

$$\partial_{\alpha\beta} f_{\gamma}(X) - \partial_{\alpha\gamma} f_{\beta}(X) = 0$$

Sp(4) field equations (Plyushchay, D.S. & Tsulaia '03) :

Fermi: $\nabla_{\alpha\beta} F_{\gamma} - \nabla_{\alpha\gamma} F_{\beta} = \frac{1}{4r} (C_{\beta(\alpha} F_{\gamma)} - C_{\gamma(\alpha} F_{\beta)})$

Bose: $\nabla_{\alpha\beta} \nabla_{\gamma\delta} B - \nabla_{\alpha\gamma} \nabla_{\beta\delta} B = \frac{1}{8r} (C_{\gamma(\delta} \nabla_{\alpha)\beta} - C_{\beta(\delta} \nabla_{\gamma)\alpha} - C_{\beta(\gamma} \nabla_{\alpha)\delta}) B + \frac{1}{32r^2} (C_{\alpha(\beta} C_{\delta)\gamma} - C_{\alpha(\gamma} C_{\delta)\beta}) B$

Generalized conformal relations between flat and Sp(4) hyperfields *(D.S. & Tsulaia '13)*

$$B(X) = \sqrt{\det G} b(X), \quad F_{\alpha}(X) = \sqrt{\det G} G_{\alpha}^{\beta} f_{\beta}(X), \quad G_{\alpha}^{\beta} = \delta_{\alpha}^{\beta} + \frac{1}{4r} X_{\alpha}^{\beta}$$

$Sp(8)$ invariant correlation functions

In flat hyperspace (*Vasiliev '01, Vasiliev & Zaikin '03*)

$$\langle b(X_1) b(X_2) \rangle = c(\det|X_1 - X_2|)^{\frac{1}{2}}$$

conformal weight

$$\langle f_\alpha(X_1) f_\beta(X_2) \rangle = c(X_1 - X_2)_{\alpha\beta}^{-1} (\det|X_1 - X_2|)^{\frac{1}{2}}$$

$$\langle b(X_1) b(X_2) b(X_3) \rangle = c(\det|X_1 - X_3|)^{-\frac{1}{4}} (\det|X_2 - X_3|)^{-\frac{1}{4}} (\det|X_1 - X_2|)^{-\frac{1}{4}}$$

In $Sp(4)$ hyperspace (*D.S. & Tsulaia '03*)

$$\langle B(X_1) B(X_2) \rangle_{Sp(4)} = \sqrt{\det G(X_1)} \sqrt{\det G(X_2)} \langle b(X_1) b(X_2) \rangle_{flat}$$

$$\langle F_\alpha(X_1) F_\beta(X_2) \rangle_{Sp(4)} = \sqrt{\det G(X_1)} \sqrt{\det G(X_2)} G_\alpha^\gamma(X_1) G_\beta^\delta(X_2) \langle f_\alpha(X_1) f_\beta(X_2) \rangle_{flat}$$

$$\langle B(X_1) B(X_2) B(X_3) \rangle_{Sp(4)} = \sqrt{\det G(X_1)} \sqrt{\det G(X_2)} \sqrt{\det G(X_3)} \langle b(X_1) b(X_2) b(X_3) \rangle_{flat}$$

$$G_\alpha^\beta(X) = \delta_\alpha^\beta + \frac{1}{4r} X_\alpha^\beta$$

Conclusion

- Free theory of the infinite number of massless HS fields in 4d flat and AdS4 space has generalized conformal $Sp(8)$ symmetry and can be compactly formulated in 10d hyperspace with the use of one scalar and one spinor field.
- Higher dimensional extension to $Sp(2M)$ invariant hyperspaces is straightforward (*Bandos, Lukierski, D.S. '99, Vasiliev '01, ...*)
known *physically relevant cases* are
 $M=2$ ($d=3$), $M=4$ ($d=4$), $M=8$ ($d=6$), $M=16$ ($d=10$)
describe conformal HS fields in corresponding space-times.
- **Important features:** hyperspace field theories possess properties of causality and locality (*Vasiliev '01*).
- Supersymmetric generalizations are available (*Bandos et. al, Vasiliev et. al., ...*)
- Hyperspace unfolded equations have Riemann's Theta-functions as solutions (*Gelfond & Vasiliev '07*)
- $Sp(8)$ hyperspace formulation of HS fields was extended to incorporate one-form gauge connections [in unfolded setting] (*Vasiliev '07*)
- **Main problem:** Whether one can construct an interacting field theory in hyperspace which would describe HS interactions in conventional space-time
 - Attempt via hyperspace SUGRA (*Bandos, Pasti, D.S., Tonin '04*)