

RICCARDO STURANI

GRAVITATIONAL WAVES
AND EFFECTIVE FIELD
THEORY METHODS TO
MODEL BINARIES

[HTTP://WWW.ICTP-SAIFR.ORG/?PAGE_ID=4372](http://www.ictp-saifr.org/?PAGE_ID=4372), VER. 2.0
DECEMBER 28th 2014

Contents

<i>Introduction</i>	7
<i>General Theory of GWs</i>	11
<i>The post-Newtonian expansion in the effective field theory approach</i>	29
<i>GW Radiation</i>	45
<i>Bibliography</i>	71

Notations

We use units $c = 1$, which means that 1 light-year (ly)=1 year $\simeq 3.16 \times 10^7$ sec = 9.46×10^{15} m.

Another useful unit is the parsec $\simeq 3.09 \times 10^{16}$ m $\simeq 3.26$ ly.

The mass of the sun is $M_\odot \simeq 1.99 \times 10^{33}$ g and its Schwarzschild radius $2G_N M_\odot \simeq 2.95$ km $\simeq 9.84 \times 10^{-6}$ sec.

We adopt the “mostly plus” ignature, i.e. the Minkowski metric is

$$\eta_{\mu\nu} = \text{diag}(-, +, +, +),$$

the Christoffel symbols is

$$\Gamma_{\mu\nu}^\rho = \frac{1}{2} g^{\rho\alpha} (g_{\alpha\mu,\nu} + g_{\alpha\nu,\mu} - g_{\mu\nu,\alpha}),$$

the Riemann tensor is

$$R_{\nu\rho\sigma}^\mu = \Gamma_{\nu\sigma,\rho}^\mu - \Gamma_{\nu\rho,\sigma}^\mu + \Gamma_{\alpha\rho}^\mu \Gamma_{\nu\sigma}^\alpha - \Gamma_{\alpha\sigma}^\mu \Gamma_{\nu\rho}^\alpha,$$

with symmetry properties

$$\begin{aligned} R_{\mu\nu\rho\sigma} &= -R_{\nu\mu\sigma\rho} = -R_{\mu\nu\sigma\rho}, \\ R_{\nu\rho\sigma}^\mu + R_{\rho\sigma\nu}^\mu + R_{\sigma\rho\nu}^\mu &= 0, \end{aligned}$$

which gives $(4 \times 3/2)^2 - 4 \times 4 = 20$ independent components,

which in $d + 1$ dimensions turn out to be $\left(\frac{(d+1)d}{2}\right)^2 - \frac{(d+1)^2 d(d-1)}{6} = \frac{(d+1)^2 d(d+2)}{12}$.

The Ricci tensor is

$$R_{\mu\nu} = R_{\mu\alpha\nu}^\alpha,$$

the Ricci scalar

$$R = R_{\mu\nu} g^{\mu\nu},$$

the Einstein tensor

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R,$$

the Weyl tensor

$$C_{\mu\nu}^{\alpha\beta} = R_{\mu\nu}^{\alpha\beta} - 2\delta_{[\mu}^{\alpha} R_{\nu]}^{\beta]} + \frac{1}{3} \delta_{[\mu}^{\alpha} \delta_{\nu]}^{\beta]} R,$$

which has 10 components in 3 + 1 dimensions or $\frac{(d+1)^2 d(d+2)}{12} - \frac{(d+1)(d+2)}{2} = (d+2)(d+1)\frac{d^2+d-6}{12}$ in $d+1$ dimensions.

Greek indices α, \dots, ω run over $d+1$ dimensions, Latin indices a, \dots, i, j, \dots over d spatial dimensions.

Fourier transform in d dimensions are defined as

$$F(\mathbf{k}) = \int d^d x F(\mathbf{x}) e^{-i\mathbf{k}\cdot\mathbf{x}},$$

$$F(\mathbf{x}) = \int \frac{d^d k}{(2\pi)^d} F(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}} = \int_{\mathbf{k}} F(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}}.$$

We will denote the modulus of a generic 3-vector \mathbf{w} by $w \equiv |\mathbf{w}|$.

Fourier transform over time is defined as

$$F(\omega) = \int dt F(t) e^{i\omega t},$$

$$F(t) = \int \frac{d\omega}{2\pi} F(\omega) e^{-i\omega t}.$$

Introduction

The existence of gravitational waves (GW) is an unavoidable prediction of General Relativity (GR): any change to a gravitating source must be communicated to distant observers no faster than the speed of light, c , leading to the existence of *gravitational radiation*, or GWs.

The more precise evidence of a system emitting GWs comes from the celebrated “Hulse-Taylor” pulsar ¹, where two neutron stars are tightly bound in a binary system with the observed decay rate of their orbit being in agreement with the GR prediction to about one part in a thousand ², see also ³ for more examples of observed GW emission from pulsar and white dwarf binary systems and sec. 6.2 of ⁴ for a pedagogical discussion.

A network of earth-based, kilometer-sized gravitational wave observatories, is currently under development: the two Laser Interferometer Gravitational-Wave Observatories (LIGO) in the US (see www.ligo.org) and the Virgo interferometer in Italy (www.virgo.infn.it) have been taking data at unprecedented sensitivities for several years and are now undergoing upgrades to their advanced stage, see e.g. ⁵ for a recent review (another smaller detector belonging to the network is the German-British Gravitational Wave Detector GEO600, www.geo600.uni-hannover.de). The gravitational detector network is planned to be joined by the Japanese KAGRA (<http://gwcenter.icrr.u-tokyo.ac.jp/en/>) detector by the end of this decade and by an additional interferometer in India (<http://www.gw-indigo.org>) by the beginning of the next decade. The advanced detector era is planned to start in the year 2015 and it is expected that few years will be necessary to reach planned sensitivity, which should allow several detections of GW events per year ⁶.

The output of such observatories is particularly sensitive to the phase Φ of GW signals and focusing on coalescing binary systems, it is possible to predict it via

$$\Phi(t) = 2 \int_{t_i}^t \omega(t') dt', \quad (1)$$

where ω is the angular velocity of the individual binary component and t_i stands for the time the signal with increasing frequency enters

¹ R A Hulse and J H Taylor. Discovery of a pulsar in a binary system. *Astrophys. J.*, 195:L51, 1975

² J. M. Weisberg and J. H. Taylor. Observations of post-newtonian timing effects in the binary pulsar psr 1913+16. *Phys. Rev. Lett.*, 52:1348, 1984

³ M. Burgay, N. D’Amico, A. Posenti, R. N. Manchester, A. G. Lyne, B. C. Joshi, M. A. McLaughlin, and M. Kramer *et al.* An increased estimate of the merger rate of double neutron stars from observations of a highly relativistic system. *Nature*, 426:531, 2003; M. Kramer and N. Wex. The double pulsar system: A unique laboratory for gravity. *Class. Quant. Grav.*, 26:073001, 2009; A. Wolszczan. A nearby 37.9-ms radio pulsar in a relativistic binary system. *Nature*, 350:688, 1991; I. H. Stairs, S. E. Thorsett, J. H. Taylor, and A. Wolszczan. Studies of the relativistic binary pulsar psr b1534+12: I. timing analysis. *Astrophys. J.*, 581:501, 2002; and J.J. Hermes, Mukremin Kilic, Warren R. Brown, D.E. Winget, Carlos Allende Prieto, et al. Rapid Orbital Decay in the 12.75-minute WD+WD Binary J0651+2844. 2012. DOI: 10.1088/2041-8205/757/2/L21

⁴ M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008

⁵ R Sturani. Ligo/virgo/geo/kagra science. In P Binétruy, G Auger, and E. Plagnol, editors, 9th LISA Symposium, Paris, volume 467 of *ASP Conference Series*, pages 293–302. Astronomical Society of the Pacific, 2012. URL <http://www.apc.univ-paris7.fr/Downloads/lisa/aspvolume.pdf>

⁶ LIGO Scientific and Virgo Collaborations. Predictions for the rates of compact binary coalescences observable by ground-based gravitational-wave detectors. *Class. Quant. Grav.*, 27:173001, 2010

the detector band-width. Note the factor 2 between the GW phase Φ and the *orbital* angular velocity ω .

Binary orbits are in general eccentric, but at the frequency we are interested in ($> 10\text{Hz}$) binary system will have circularized, see sec. 4.1.3 of ⁷ for a quantitative analysis of orbit circularization. For circular orbit the binding energy can be expressed in terms of a single parameter, say the relative velocity of the binary system components v , which by the virial theorem $v^2 \simeq G_N M/r$, being G_N the standard Newton constant, M the total mass of the binary system, and r the orbital separation between its constituents. Note that the virial relationship, or its equivalent (on circular orbits) Kepler law $\omega^2 \simeq G_N M/r^3$, are not exact in GR, but only at Newtonian level.

At Newtonian level (for spin-less binary constituents) one has for the energy E of circular orbits:

$$E(v) = -\frac{1}{2}\eta Mv^2 \left(1 + e_{v^2}(\eta)v^2 + e_{v^4}(\eta)v^4 + \dots\right), \quad (2)$$

where $\eta \equiv m_1 m_2 / M^2$ is the symmetric mass ratio and the $e_{v^n}(\eta)$ coefficients stand for GR corrections to the Newtonian formula, and only even power of v are involved for the conservative Energy. For the radiated flux $F(v)$ the leading term is the Einstein quadrupole formula, which we will derive in sec. , that in the circular orbit case reduces to

$$F(v) = \frac{32\eta^2}{5G_N} v^{10} \left(1 + f_{v^2}(\eta)v^2 + f_{v^3}(\eta)v^3 + \dots\right). \quad (3)$$

Note that using $v = \omega r$ together with the virial relation we have

$$v = \omega r \implies \omega \simeq \frac{1}{G_N M} v^3, \quad (4)$$

allowing to re-write (note that during the coalescence v increases monotonically) eq. (1) as

$$\Phi(v) \simeq \frac{2}{G_N M} \int_{v_i}^v v^3 \frac{dE/dv}{dE/dt} dv = \frac{5}{16\eta} \int_{v_i}^v \frac{1}{v^6} \left(1 + p_{v^2}v^2 + p_{v^3}v^3 + \dots\right) dv, \quad (5)$$

where we used $dE/dt = -F$. Since the phase has to be matched with $O(1)$ precision, corrections at least $O(v^6)$ must be considered.

The aim of this course is to show how to compute the E, F functions at required perturbative order. We thus have to treat the binary problem perturbatively, the actual expansion parameter will be $(G_N M \pi f_{GW}) = (G_N M \omega)^{1/3} \simeq v$, which represents an expansion around the Minkowski space. Such perturbative expansion of GR has been proven very useful to treat the binary problem and it goes under the name of post-Newtonian (PN) expansion to GR.

The approach to solving for the dynamics of the two body problem adopted here relies on an *effective field theory methods*, originally

⁷ M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008

proposed in ⁸. The two body problem is a system which exhibits a clear separation of scales: the size of the compact objects r_s , like black holes and/or neutron stars, the orbital separation r and the gravitational wave-length λ . Using again the virial theorem the hierarchy $r_s < r \sim r_s/v^2 < \lambda \sim r/v$ can be established.

The author of the present notes recommends the following reviews: for a review of the PN theory see ⁹, for a pedagogical book on GWs see ¹⁰, for an astrophysics oriented review on GWs see ¹¹, for a data-analysis oriented review see ¹², for reviews on effective field theory methods for GR see ¹³ and ¹⁴.

These are the notes of the course held at ICTP-SAIFR in August 2013 and they are not meant in any way to replace or improve the extensive literature existent on the topic but rather to collect in single document the material relevant for this course, which could otherwise be found scattered in different places.

⁸ Walter D. Goldberger and Ira Z. Rothstein. An Effective field theory of gravity for extended objects. *Phys.Rev.*, D73:104029, 2006

⁹ Luc Blanchet. Gravitational radiation from post-newtonian sources and inspiralling compact binaries. *Living Reviews in Relativity*, 9(4), 2006. URL <http://www.livingreviews.org/lrr-2006-4>

¹⁰ M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008

¹¹ B.S. Sathyaprakash and B.F. Schutz. Physics, Astrophysics and Cosmology with Gravitational Waves. *Living Rev.Rel.*, 12:2, 2009

¹² Alessandra Buonanno. Gravitational waves. 2007. URL <http://arxiv.org/abs/arXiv:0709.4682>

¹³ W. D. Goldberger. Les houches lectures on effective field theories and gravitational radiation. In *Les Houches Summer School - Session 86: Particle Physics and Cosmology: The Fabric of Spacetime*, 2007

¹⁴ Stefano Foffa and Riccardo Sturani. Effective field theory methods to model compact binaries. *Class.Quant.Grav.*, 31(4):043001, 2014. DOI: 10.1088/0264-9381/31/4/043001

General Theory of GWs

Expansion around Minkowski

We start by recalling the Einstein equations

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi G_N T_{\mu\nu}, \quad (6)$$

however it will be useful for our purposes to work also at the level of the action

$$S_{EH} = \frac{1}{16\pi G_N} \int dt d^d x \sqrt{-g} R, \quad (7)$$

$$S_m \rightarrow \delta S_m = \frac{1}{2} \int dt d^d x \sqrt{-g} T^{\mu\nu} \delta g_{\mu\nu}. \quad (8)$$

Here we focus on an expansion around the Minkowski space-time

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} \quad |h_{\mu\nu}| \ll 1, \quad (9)$$

and we are interested in a systematic expansion in powers of $h_{\mu\nu}$. As in the binary system case the metric perturbation $|h_{\mu\nu}| \sim G_N m/r \sim v^2$, a suitable velocity expansion will have to be considered.

GR admits invariance under general coordinate transformations

$$x^\mu \rightarrow x'^\mu = x^\mu + \xi^\mu(x), \quad (10)$$

which change the metric according to

$$\begin{aligned} g_{\mu\nu}(x) \rightarrow g'_{\mu\nu}(x') &= g_{\rho\sigma}(x) \frac{\partial x^\rho}{\partial x'^\mu} \frac{\partial x^\sigma}{\partial x'^\nu} \\ h_{\mu\nu}(x) \rightarrow h'_{\mu\nu}(x') &= h_{\mu\nu}(x) - (\partial_\mu \xi_\nu + \partial_\nu \xi_\mu). \end{aligned} \quad (11)$$

At linear order around a Minkowski background we have ($h \equiv \eta_{\mu\nu} h^{\mu\nu}$)

$$\begin{aligned} R_{\mu\nu\rho\sigma} &= \frac{1}{2} (\partial_\nu \partial_\rho h_{\mu\sigma} + \partial_\mu \partial_\sigma h_{\nu\rho} - \partial_\mu \partial_\rho h_{\nu\sigma} - \partial_\nu \partial_\sigma h_{\mu\rho}), \\ R_{\mu\nu} &= \frac{1}{2} (\partial_\rho \partial_\mu h_\nu^\rho + \partial_\rho \partial_\nu h_\mu^\rho - \square h_{\mu\nu} - \partial_\mu \partial_\nu h), \\ R &= \partial_\mu \partial_\nu h^{\mu\nu} - \square h, \\ G_{\mu\nu} &= \frac{1}{2} (\partial_\rho \partial_\mu h_\nu^\rho + \partial_\rho \partial_\nu h_\mu^\rho - \square h_{\mu\nu} - \partial_\mu \partial_\nu h - \eta_{\mu\nu} \partial_\rho \partial_\sigma h^{\rho\sigma} + \eta_{\mu\nu} \square h), \end{aligned} \quad (12)$$

and we remind that with the metric signature adopted here $\square = \partial_i \partial^i - \partial_t^2$.

The formula for $G_{\mu\nu}$ can be used to write the Einstein equations as

$$\square \bar{h}_{\mu\nu} + \eta_{\mu\nu} \partial_\rho \partial_\sigma \bar{h}^{\rho\sigma} - \partial_\mu \partial_\rho \bar{h}_\nu^\rho - \partial_\nu \partial_\rho \bar{h}_\mu^\rho = -16\pi G_N T_{\mu\nu} \quad (13)$$

whre $\bar{h}_{\mu\nu} \equiv h_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} h$ which transforms as

$$\bar{h}_{\mu\nu} \rightarrow \bar{h}'_{\mu\nu} = \bar{h}_{\mu\nu} - \xi_{\mu,\nu} - \xi_{\nu,\mu} + \eta_{\mu\nu} \xi^\alpha{}_\alpha.$$

Thanks to the gauge freedom it is always possible to go to a *class* of gauges specified by the *Lorentz condition*

$$\partial^\mu \bar{h}_{\mu\nu} = 0, \quad (14)$$

for instance by choosing $\square \xi_\mu = \partial^\nu \bar{h}_{\mu\nu}$ we can impose $\partial^\mu \bar{h}'_{\mu\nu} = 0$. Note that in this way we have not completely fixed the gauge as we can always perform a transformation with $\square \xi_\mu = 0$ and still remain in the Lorentz gauge class, in which the equation of motions assume the very simple form

$$\square \bar{h}_{\mu\nu} = -16\pi G_N T_{\mu\nu}, \quad (15)$$

showing that *in this class of gauges* all gravity components satisfy a wave-like equation. However we will see later in this section that working with gauge invariant variables (though non-local) shows that only 2 degrees of freedom are physical and *radiative*, 4 more are physical and *non-radiative* and finally 4 are pure gauge, and can be killed by the Lorentz condition in eq. (14), for instance.

It will be useful to solve eq. (15) by using the Green function G with appropriate boundary conditions, where the G is defined as

$$\square_x G(x-y) = \delta^{(4)}(x-y). \quad (16)$$

The explicit form of the Green functions with time-retarded and time-advanced boundary conditions are in the direct space (see e.g. sec. 6.4 of ¹⁵, with a pre-factor different by -4π because of a different definition in eq. (16)

$$\begin{aligned} G_{ret}(t, \mathbf{x}) &= -\delta(t-r) \frac{1}{4\pi r}, \\ G_{adv}(t, \mathbf{x}) &= -\delta(t+r) \frac{1}{4\pi r}, \end{aligned} \quad (17)$$

where $r \equiv |\mathbf{x}| > 0$ (note that $G_{ret}(t, \mathbf{x}) = G_{adv}(-t, \mathbf{x})$). By solving ex. 3,4 one can show that these are indeed Green functions for the eq. (16). For the Feynman prescription of the Green function G_F see ex. 5, where the motivated student is asked to demonstrate that the G_F ensures pure incoming wave at past infinity and pure outgoing wave at future infinity.

¹⁵ John David Jackson. *Classical Electrodynamics*. John Wiley & Sons, iii edition, 1999

Let us now focus on the wave eq. (15) in the vacuum case

$$\square \bar{h}_{\mu\nu} = 0 \quad (18)$$

and define

$$\zeta_{\mu\nu} = \partial_\nu \zeta_\mu + \partial_\mu \zeta_\nu - \eta_{\mu\nu} \partial^\alpha \zeta_\alpha. \quad (19)$$

Now $\square \zeta_\mu = 0 \implies \square \zeta_{\mu\nu} = 0$, with $\delta \bar{h}_{\mu\nu} = \zeta_{\mu\nu}$. Since now the free wave equation is satisfied by both the gravitational field $\bar{h}_{\mu\nu}$ and by the residual gauge transformation parametrized by ζ_μ that preserves the Lorentz condition, we can use the four available ζ_μ to set four conditions on $\bar{h}_{\mu\nu}$. In particular ζ_0 can be used to make h vanish (so that $\bar{h}_{\mu\nu} = h_{\mu\nu}$) and the three ζ_i can be used to make the three \bar{h}_{0i} vanish. The Lorentz condition eq. (14) for $\mu = 0$ will now look like $\partial^0 h_{00} = 0$, which means that h_{00} is constant in time, hence not contributing to any GW. In conclusion, in vacuum one can set

$$h_{0\mu} = h = \partial^i h_{ij} = 0, \quad (20)$$

defining the *transverse traceless*, or TT gauge, which then describe only the physical GW propagating in vacuum. For a wave propagating along the $\mu = 3$ axis, for instance, the wave eq. (15) admits the solution $h_{\mu\nu}(t - z)$ and the gauge condition $\partial^j h_{ij} = 0$ reads $h_{i3} = 0$. In terms of the tensor components we have

$$h_{\mu\nu}^{(TT)} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & h_+ & h_\times & 0 \\ 0 & h_\times & -h_+ & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (21)$$

Given a plane wave solution $h_{kl}^{(GW)}$ propagating in the generic \hat{n} direction outside the source, which is in the Lorentz gauge, but not yet TT-ed, its TT form can be obtained by applying the projector $\Lambda_{ij,kl}$ defined as

$$\begin{aligned} \Lambda_{ij,kl}(\hat{n}) &= \frac{1}{2} \left[P_{ik}P_{jl} + P_{il}P_{jk} - P_{ij}P_{kl} \right], \\ P_{ij}(\hat{n}) &= \delta_{ij} - n_i n_j, \end{aligned} \quad (22)$$

according to

$$h_{ij}^{(TT-GW)} = \Lambda_{ij,kl} h_{kl}^{(GW)}. \quad (23)$$

The Λ projector ensures transversality and tracelessness of the resulting tensor (starting from a tensor in the Lorentz gauge!).

The TT gauge cannot be imposed there where $T_{\mu\nu} \neq 0$, as we cannot set to 0 any component of $\bar{h}_{\mu\nu}$ which satisfies a $\square \bar{h}_{\mu\nu} \neq 0$ equation by using a ζ_μ which satisfies a $\square \zeta_\mu = 0$ equation (and hence $\square \zeta_{\mu\nu} = 0$).

Radiative degrees of freedom of $h_{\mu\nu}$

Following sec. 2.2 of ¹⁶ we show that the gravitational perturbation $h_{\mu\nu}$ has 6 physical degrees of freedom: 4 constrained plus 2 radiative. The argument is based on a Minkowski equivalent of Bardeen's gauge-invariant cosmological perturbation formalism.

We begin by defining the decomposition of the metric perturbation $h_{\mu\nu}$, in any gauge, into a number of irreducible pieces. Assuming that $h_{\mu\nu} \rightarrow 0$ as $r \rightarrow \infty$, we decompose $h_{\mu\nu}$ into a number of irreducible quantities ϕ , β_i , γ , H , ε_i , λ and $h_{ij}^{(\text{TT})}$ via the equations

$$h_{00} = 2\phi, \quad (24)$$

$$h_{0i} = \beta_i + \partial_i \gamma, \quad (25)$$

$$h_{ij} = h_{ij}^{(\text{TT})} + \frac{1}{3}H\delta_{ij} + \partial_{(i}\varepsilon_{j)} + \left(\partial_i\partial_j - \frac{1}{3}\delta_{ij}\nabla^2\right)\lambda, \quad (26)$$

together with the constraints

$$\partial_i\beta_i = 0 \quad (1 \text{ constraint}) \quad (27)$$

$$\partial_i\varepsilon_i = 0 \quad (1 \text{ constraint}) \quad (28)$$

$$\partial_i h_{ij}^{(\text{TT})} = 0 \quad (3 \text{ constraints}) \quad (29)$$

$$\delta^{ij}h_{ij}^{(\text{TT})} = 0 \quad (1 \text{ constraint}) \quad (30)$$

and boundary conditions

$$\gamma \rightarrow 0, \quad \varepsilon_i \rightarrow 0, \quad \lambda \rightarrow 0, \quad \nabla^2\lambda \rightarrow 0 \quad (31)$$

as $r \rightarrow \infty$. Here $H \equiv \delta^{ij}h_{ij}$ is the trace of the *spatial* portion of the metric perturbation. The spatial tensor $h_{ij}^{(\text{TT})}$ is transverse and traceless, and is the TT piece of the metric discussed above which contains the physical radiative degrees of freedom. The quantities β_i and $\partial_i\gamma$ are the transverse and longitudinal pieces of h_{ti} . The uniqueness of this decomposition follows from taking a divergence of Eq. (25) giving $\nabla^2\gamma = \partial_i h_{ti}$, which has a unique solution by the boundary condition (31). Similarly, taking two derivatives of Eq. (26) yields the equation $2\nabla^2\nabla^2\lambda = 3\partial_i\partial_j h_{ij} - \nabla^2 H$, which has a unique solution by Eq. (31). Having solved for λ , one can obtain a unique ε_i by solving $3\nabla^2\varepsilon_i = 6\partial_j h_{ij} - 2\partial_i H - 4\partial_i\nabla^2\lambda$.

The total number of free functions in the parameterization (24) – (26) of the metric is 16: 4 scalars (ϕ , γ , H , and λ), 6 vector components (β_i and ε_i), and 6 symmetric tensor components ($h_{ij}^{(\text{TT})}$). The number of constraints (27) – (30) is 6, so the number of independent variables in the parameterization is 10, consistent with a symmetric 4×4 tensor.

We next discuss how the variables ϕ , β_i , γ , H , ε_i , λ and $h_{ij}^{(\text{TT})}$ transform under gauge transformations ζ^a with $\zeta^a \rightarrow 0$ as $r \rightarrow \infty$. We

¹⁶ Eanna E. Flanagan and Scott A. Hughes. The basics of gravitational wave theory. *New J. Phys.*, 7:204, 2005

parameterize such gauge transformation as

$$\xi_\mu = (\xi_t, \xi_i) \equiv (A, B_i + \partial_i C), \quad (32)$$

where $\partial_i B_i = 0$ and $C \rightarrow 0$ as $r \rightarrow \infty$; thus B_i and $\partial_i C$ are the transverse and longitudinal pieces of the spatial gauge transformation.

Decomposing this transformed metric into its irreducible pieces yields the transformation laws

$$\phi \rightarrow \phi - \dot{A}, \quad (33)$$

$$\beta_i \rightarrow \beta_i - \dot{B}_i, \quad (34)$$

$$\gamma \rightarrow \gamma - A - \dot{C}, \quad (35)$$

$$H \rightarrow H - 2\nabla^2 C, \quad (36)$$

$$\lambda \rightarrow \lambda - 2C, \quad (37)$$

$$\varepsilon_i \rightarrow \varepsilon_i - 2B_i, \quad (38)$$

$$h_{ij}^{(\text{TT})} \rightarrow h_{ij}^{(\text{TT})}. \quad (39)$$

Gathering terms, we see that the following combinations of these functions are gauge invariant:

$$\Phi \equiv -\phi + \dot{\gamma} - \frac{1}{2}\ddot{\lambda}, \quad (40)$$

$$\Theta \equiv \frac{1}{3} \left(H - \nabla^2 \lambda \right), \quad (41)$$

$$\Xi_i \equiv \beta_i - \frac{1}{2}\dot{\varepsilon}_i; \quad (42)$$

$h_{ij}^{(\text{TT})}$ is gauge-invariant without any further manipulation. In the Newtonian limit Φ reduces to the Newtonian potential Φ_N , while $\Theta = -2\Phi_N$. The total number of free, gauge-invariant functions is 6: 1 function Θ ; 1 function Φ ; 3 functions Ξ_i , minus 1 due to the constraint $\partial_i \Xi_i = 0$; and 6 functions $h_{ij}^{(\text{TT})}$, minus 3 due to the constraints $\partial_i h_{ij}^{(\text{TT})} = 0$, minus 1 due to the constraint $\delta^{ij} h_{ij}^{(\text{TT})} = 0$. This is in keeping with the fact that in general the 10 metric functions contain 6 physical and 4 gauge degrees of freedom.

We would now like to enforce Einstein's equation. Before doing so, it is useful to first decompose the stress energy tensor in a manner similar to that of our decomposition of the metric. We define the quantities ρ , S_i , S , P , σ_{ij} , σ_i and σ via the equations

$$T_{00} = \rho, \quad (43)$$

$$T_{0i} = S_i + \partial_i S, \quad (44)$$

$$T_{ij} = P\delta_{ij} + \sigma_{ij} + \partial_{(i}\sigma_{j)} + \left(\partial_i \partial_j - \frac{1}{3}\delta_{ij}\nabla^2 \right) \sigma, \quad (45)$$

together with the constraints

$$\partial_i S_i = 0, \quad (46)$$

$$\partial_i \sigma_i = 0, \quad (47)$$

$$\partial_i \sigma_{ij} = 0, \quad (48)$$

$$\delta^{ij} \sigma_{ij} = 0, \quad (49)$$

and boundary conditions

$$S \rightarrow 0, \quad \sigma_i \rightarrow 0, \quad \sigma \rightarrow 0, \quad \nabla^2 \sigma \rightarrow 0 \quad (50)$$

as $r \rightarrow \infty$. These quantities are not all independent. The variables ρ , P , S_i and σ_{ij} can be specified arbitrarily; stress-energy conservation ($\partial^a T_{ab} = 0$) then determines the remaining variables S , σ , and σ_i via

$$\nabla^2 S = \dot{\rho}, \quad (51)$$

$$\nabla^2 \sigma = -\frac{3}{2}P + \frac{3}{2}\dot{S}, \quad (52)$$

$$\nabla^2 \sigma_i = 2\dot{S}_i. \quad (53)$$

We now compute the Einstein tensor from the metric (24) – (26). The result can be expressed in terms of the gauge invariant observables:

$$G_{00} = -\nabla^2 \Theta, \quad (54)$$

$$G_{0i} = -\frac{1}{2}\nabla^2 \Xi_i - \partial_i \dot{\Theta}, \quad (55)$$

$$G_{ij} = -\frac{1}{2}\square h_{ij}^{(\text{TT})} - \partial_{(i} \dot{\Xi}_{j)} - \frac{1}{2}\partial_i \partial_j (2\Phi + \Theta) + \delta_{ij} \left[\frac{1}{2}\nabla^2 (2\Phi + \Theta) - \ddot{\Theta} \right]. \quad (56)$$

We finally enforce Einstein's equation $G_{\mu\nu} = 8\pi T_{\mu\nu}$ and simplify using the conservation relations (51) – (53); this leads to the following field equations:

$$\nabla^2 \Theta = -8\pi\rho, \quad (57)$$

$$\nabla^2 \Phi = 4\pi(\rho + 3P - 3\dot{S}), \quad (58)$$

$$\nabla^2 \Xi_i = -16\pi S_i, \quad (59)$$

$$\square h_{ij}^{(\text{TT})} = -16\pi\sigma_{ij}. \quad (60)$$

Notice that **only the metric components $h_{ij}^{(\text{TT})}$ obey a wave-like equation**. The other variables Θ , Φ and Ξ_i are determined by Poisson-type equations. Indeed, in a purely vacuum spacetime, the

field equations reduce to five Laplace equations and a wave equation:

$$\nabla^2 \Theta^{\text{vac}} = 0, \quad (61)$$

$$\nabla^2 \Phi^{\text{vac}} = 0, \quad (62)$$

$$\nabla^2 \Xi_i^{\text{vac}} = 0, \quad (63)$$

$$\square h_{ij}^{(\text{TT}),\text{vac}} = 0. \quad (64)$$

This manifestly demonstrates that only the $h_{ij}^{(\text{TT})}$ metric components — the transverse, traceless degrees of freedom of the metric perturbation — characterize the radiative degrees of freedom in the spacetime. Although it is possible to pick a gauge in which other metric components *appear* to be radiative, they will not be: Their radiative character is an illusion arising due to the choice of gauge or coordinates.

The field equations (57) – (60) also demonstrate that, far from a dynamic, radiating source, the time-varying portion of the physical degrees of freedom in the metric is dominated by $h_{ij}^{(\text{TT})}$. If we expand the gauge invariant fields Φ , Θ , Ξ_i and $h_{ij}^{(\text{TT})}$ in powers of $1/r$, then, at sufficiently large distances, the leading-order $O(1/r)$ terms will dominate. For the fields Θ , Φ and Ξ_i , the coefficients of the $1/r$ pieces are combinations of the conserved quantities given by the mass $\int d^3x T_{00}$, the linear momentum $\int d^3x T_{0i}$ and the angular momentum $\int d^3x (x_i T_{0j} - x_j T_{0i})$. Thus, the only time-varying piece of the physical degrees of freedom in the metric perturbation at order $O(1/r)$ is the TT piece $h_{ij}^{(\text{TT})}$.

Although the variables Φ , Θ , Ξ_i and $h_{ij}^{(\text{TT})}$ have the advantage of being gauge invariant, they have the disadvantage of being *non-local* (a part from $h_{ij}^{(\text{TT})}$ computation of these variables at a point requires knowledge of the metric perturbation $h_{\mu\nu}$ everywhere). This non-locality obscures the fact that the physical, non-radiative degrees of freedom are causal, a fact which is explicit in Lorentz gauge. One way to see that the gauge invariant degrees of freedom are causal is to combine the vacuum wave equation eq. (18) for the metric perturbation with the expression (12) for the gauge-invariant Riemann tensor. This gives the wave equation $\square R_{\alpha\beta\gamma\delta} = 0$. Moreover, many observations that seek to detect GWs are sensitive only to the value of the Riemann tensor at a given point in space (see sec.). For example, the Riemann tensor components R_{itjt} are given in terms of the gauge invariant variables as

$$R_{i0j0} = -\frac{1}{2} \ddot{h}_{ij}^{(\text{TT})} + \Phi_{,ij} + \dot{\Xi}_{(i,j)} - \frac{1}{2} \ddot{\Theta} \delta_{ij}. \quad (65)$$

Thus, at least certain combinations of the gauge invariant variables are locally observable.

Energy of GWs

We have shown that only the TT part of the metric is actually a radiative degree of freedom, i.e. a GW, which is capable of transporting energy, momentum and angular momentum from the source emitting them. In order to derive the expression for such quantities we follow here sec. 1.4 of ¹⁷.

In principle it is not unambiguous to separate the background metric from the perturbation, but a natural splitting between space-time background and GWs arise when there is a clear separation of scales: if the variation scale of $h_{\mu\nu}$ is λ and the variation scale of the background is $L_B \ll \lambda$ a separation is possible. We can e.g. average over a time scale $\bar{t} \gg \lambda$ and obtain

$$\bar{R}_{\mu\nu} = 8\pi G_N \left(\tau_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} \tau \right) - \langle R^{(2)} \rangle, \quad (66)$$

where $\bar{R}_{\mu\nu}$ is the Ricci tensor computed on the background metric (and then vanishing in an expansion over Minkowski background), and $R_{\mu\nu}^{(2)}$ is the part of the Ricci tensor quadratic in the GW perturbation (no part linear in the perturbation survives after averaging). The $-\langle R^{(2)} \rangle$ in the above equation can be interpreted as giving contribution to an effective energy momentum tensor of the GWs $\tau_{\mu\nu}$ given by

$$\tau_{\mu\nu} = -\frac{1}{8\pi G_N} \langle R_{\mu\nu}^{(2)} - \frac{1}{2} \eta_{\mu\nu} R^{(2)} \rangle \quad (67)$$

(and $\tau = \langle R^{(2)} \rangle / (8\pi G_N)$). The expression for $R_{\mu\nu}^{(2)}$ is quite lengthy, see eq. (1.131) of ¹⁸, but using the Lorentz condition, $h = 0$ and neglecting terms which vanish on the equation of motion $\square \bar{h}_{\mu\nu} = 0$ we have

$$\langle R_{\mu\nu}^{(2)} \rangle = -\frac{1}{4} \langle \partial_\mu h_{\alpha\beta} h^{\alpha\beta} \partial_\nu \rangle, \quad (68)$$

$$\tau_{\mu\nu} = \frac{1}{32\pi G_N} \langle \partial_\mu h_{\alpha\beta} \partial_\nu h^{\alpha\beta} \rangle, \quad (69)$$

This effective energy-momentum tensor is gauge invariant and thus depend only on $h_{ij}^{(TT)}$, giving

$$\tau^{00} = \frac{1}{16\pi G_N} \langle \dot{h}_+^2 + \dot{h}_\times^2 \rangle$$

For a plane wave travelling along the z direction we have $\tau_{01} = 0 = \tau_{02}$ and $\partial_z h_{ij}^{(TT)} = \partial^0 h_{ij}^{(TT)}$ and then $\tau^{03} = \tau^{00}$. For a spherical wave $\partial_r h_{ij}^{(TT)} = \partial^0 h_{ij}^{(TT)} + O(1/r^2)$, so similarly $\tau^{0r} = \tau^{00}$.

¹⁷ M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008

¹⁸ M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008

The time derivative of the GW energy E_V (or energy flux dE_V/dt) can be written as

$$\begin{aligned} \frac{dE_V}{dt} &= \int_V d^3x \partial_0 \tau^{00} \\ &= \int_V d^3x \partial_i \tau^{0i} \\ &= \int_S dA n_i \tau^{0i} = \\ &= \int_S dA \tau^{00} \end{aligned} \quad (70)$$

The outward propagating GW carries then an energy flux $F = \frac{dE}{dt}$

$$F = \frac{r^2}{32\pi G_N} \int d\Omega \langle \dot{h}_{ij}^{(TT)} \dot{h}_{ij}^{(TT)} \rangle \quad (71)$$

or equivalently

$$F = \frac{1}{16\pi G_N} \langle \dot{h}_+^2 + \dot{h}_\times^2 \rangle. \quad (72)$$

The linear momentum P_V^i of GW is

$$P_V^k = \int d^3x \tau^{0k}. \quad (73)$$

Considering a GW propagating radially outward, we have

$$\dot{P}_V^i = - \int_S \tau^{0i}, \quad (74)$$

hence the momentum carried away by the outward-propagating wave is $\frac{dP^i}{d\lambda dt} = \dot{t}^{0i}$. In terms of the GW amplitude it is

$$\frac{dP^i}{dt} = - \frac{r^2}{32\pi G_N} \int d\Omega \langle \dot{h}_{jk}^{(TT)} \partial^i h_{jk}^{(TT)} \rangle. \quad (75)$$

Note that, if τ^{0k} is odd under a parity transformation $\mathbf{x} \rightarrow -\mathbf{x}$, then the angular integral of eq. (75) vanishes. For the angular momentum of GW we refer to sec. 2.1.3 of ¹⁹.

¹⁹ M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008

Interaction of GWs with interferometric detectors

We now describe the interaction of a GW with a simplified experimental apparatus, following the discussion of sec. 1.3.3 of ²⁰. In a typical interferometer lights goes back and forth in orthogonal arms and recombining the photons after different trajectories very precise length measurements can be performed.

The dynamics of a massive point particle can be inferred from the action

$$S = -m \int d\tau = \int \sqrt{-g_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}} d\lambda, \quad (76)$$

²⁰ M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008

whose variation with respect to the particle trajectory x^μ gives the geodesic equation of motion

$$\frac{d^2 x^\mu}{d\tau^2} + \frac{1}{2} g^{\mu\alpha} \left(g_{\alpha\nu,\rho} + g_{\alpha\rho,\nu} - \frac{1}{2} g_{\nu\rho,\alpha} \right) \frac{dx^\nu}{d\tau} \frac{dx^\rho}{d\tau}. \quad (77)$$

Let us apply the above equation to the motion of a mirror, at rest at $\tau = 0$ in the position $\mathbf{x}_m = (L, 0, 0)$, in a laser interferometer under the influence of a GW propagating along the z direction:

$$\left. \frac{d^2 x^i}{d\tau^2} \right|_{\tau=0} = - \Gamma_{00}^i \left(\left. \frac{dx^0}{d\tau} \right|_{\tau=0} \right)^2. \quad (78)$$

In the TT gauge $\Gamma_{00}^i = 0$, showing that an object initially at rest will remain at rest even during the passage of a GW. This does not mean that the GW will have no effect, as the *physical* distance l between the mirror and the beam splitter, say, at $\mathbf{x}_{bs} = (0, 0, 0)$ is given by

$$l = \int_0^L \sqrt{g_{xx}} dx \simeq \left(1 + \frac{1}{2} h_+(t) \right) L,$$

which shows how physical relative distance change with time (we have assumed that h_+ does not depend on x , which is correct for $\lambda_{GW} \gg L$).

It is instructive to review this derivation in a different frame, the *proper detector frame*, whose coordinates allow a more transparent interpretation, as all physical results are *independent of frame choices*. Experimentally one has the mirror and the beam splitter, which are “freely-falling”, as they are suspended to the ceiling forming a pendulum with very little friction and low typical frequency (\sim few Hz), meaning that over frequency scale $10 - 10^3$ Hz they behave as freely-falling particles, see fig. 1. Standard rulers on the other hand are made of tightly bound objects, endowed with friction and restoring forces. The distance they measure $\xi(t)$ under the influence of a monochromatic wave frequency ω undergoes oscillation satisfying the equation

$$\ddot{\xi}(t) + \gamma \omega_0 \dot{\xi}(t) + \omega_0^2 \xi(t) = -\frac{\omega^2}{2} h_0 \cos(\omega t) L \quad (79)$$

with solution

$$\xi(t) = \frac{1}{2} L h_0 \omega^2 \frac{(\omega^2 - \omega_0^2) \cos(\omega t) - \gamma \omega_0 \omega \sin(\omega t)}{(\omega^2 - \omega_0^2)^2 + \gamma^2 \omega_0^2 \omega^2}, \quad (80)$$

showing that for $\omega \gg \omega_0$ the mirror is indeed freely-falling, as it follows the GW time behaviour.

In the proper detector frame, which is the freely falling frame for the observer at the origin at the coordinates, a general metric can be

written as

$$ds^2 \simeq -dt^2 \left(1 + R_{0i0j}x^i x^j + \dots\right) - 2dtdx^i \left(\frac{2}{3}R_{0ijk}x^j x^k + \dots\right) + dx^i dx^j \left(\delta_{ij} - \frac{1}{3}R_{ijkl}x^k x^l \dots\right) \quad (81)$$

where terms of cubic and higher order in x have been omitted. The trajectory $x_0^\mu(\tau) = \delta^{\mu 0}\tau$ is clearly a geodesic, and we can consider the *geodesic deviation* equation, which gives the time evolution separation of two nearby geodesics $x_0^\mu(\tau)$ and $x_0^\mu(\tau) + \zeta^\mu(\tau)$

$$\frac{D^2 \zeta^i}{d\tau^2} = -R^i{}_{0j0} \zeta^j \left(\frac{dx^0}{d\tau}\right)^2. \quad (82)$$

In the proper detector frame, the measure of coordinate distances with respect to the origin gives actually a measure of physical distance (up to terms $\sim hL^3/\lambda^3$). The eq.(82) can be recast into

$$\ddot{\zeta} = \frac{1}{2}\ddot{h}_{ij}\zeta^j, \quad (83)$$

where an overdot stands for a derivative with respect to t and terms of order h^2 have been neglected. In the proper detector frame, the effect of GWs on a point particle of mass m placed at coordinate ζ can be described in terms of a Newtonian force F_i

$$F_i = \frac{m}{2}\ddot{h}_{ij}^{(TT)}(t)\zeta^j \quad (84)$$

(we neglect again the space dependence of h_+ as typically GW wavelength $\lambda \gg L$).




Figure 1: Interferometer scheme. Light emitted from the laser is shared by the two orthogonal arms after going through the beam splitter. After bouncing at the end mirrors it is recombined at the photo-detector.

The laser light travels in two orthogonal arms of the interferometer and the electric fields are finally recombined on the photo-detector. The reflection off a 50-50 beam splitter can be modeled by multiplying the amplitude of the incoming field by $1/\sqrt{2}$ for reflection on one side and $-1/\sqrt{2}$ for reflection on the other, while transmission multiplies it by $1/\sqrt{2}$ and reflection by the end mirrors by -1 , see sec. 2.4.1 of ²¹ for details.

²¹ Andreas Freise and Kenneth A. Strain. Interferometer techniques for gravitational-wave detection. *Living Reviews in Relativity*, 13(1), 2010. DOI: 10.12942/lrr-2010-1. URL <http://www.livingreviews.org/lrr-2010-1>

Setting the mirrors at positions $(L_x, 0, 0)$ and $(0, L_y, 0)$ and the beam splitter at the origin of the coordinates, we can compute the phase change of the laser electric field moving in the x and y cavity which are eventually recombined on the photo-multiplier. For the light propagating along the x -axis, using the TT metric (21) for a z -propagating GW, one has

$$\begin{aligned} L_x &= (t_{1x} - t_{0x}) - \frac{1}{2} \int_{t_{0x}}^{t_{1x}} h_+(t') dt' , \\ L_x &= (t_{2x} - t_{1x}) - \frac{1}{2} \int_{t_{1x}}^{t_{2x}} h_+(t') dt' , \end{aligned} \quad (85)$$

where $t_{0,1,2}$ stand respectively for the time when the laser leaves the beam splitter, bounces off the mirror, returns to the beam splitter. The time at which laser from the two arms is combined at the beam splitter is common: $t_{2x} = t_{2y} = t_2$, and using $h_+(t) = h_0 \cos(\omega_{GW}t)$ we get

$$\begin{aligned} t_2 &= t_{0x} + 2L_x + \frac{h_0}{2\omega_{GW}} [\sin(\omega_{GW}(t_0 + 2L)) - \sin(\omega_{GW}t_0)] \\ &= t_{0x} + 2L_x + h_0 L_x \frac{\sin(\omega_{GW}L_x)}{\omega_{GW}L_x} \cos(\omega_{GW}(t_0 + 2L_x)) \end{aligned} \quad (86)$$

where the trigonometric identity $\sin(\alpha + 2\beta) = \sin(\alpha) + 2\cos(\alpha + \beta)\sin\beta$ has been used. Using that the phase of the laser field x (y) at recombination time t_2 is the same it had at the time it left the beam splitter at time t_{0x} (t_{0y}), we can write

$$\begin{aligned} E^{(x)}(t_2) &= -\frac{1}{2} E_0 e^{-i\omega_L t_{0x}} = \\ &= -\frac{1}{2} E_0 e^{-i\omega_L(t_2 - 2L) + i\phi_0 + i\Delta\phi_x} \end{aligned} \quad (87)$$

where

$$\begin{aligned} L &\equiv \frac{L_x + L_y}{2} \\ \phi_0 &= \omega_L(L_x - L_y) \\ \Delta\phi_x &= h_0 \omega_L L \frac{\sin(\omega_{GW}L)}{\omega_{GW}L} \cos(\omega_{GW}(t - L)). \end{aligned} \quad (88)$$

where in the terms $O(h)$ we have set $L_x \simeq L_y \simeq L$. Analogously for the field that traveled through the y - arm

$$\begin{aligned} E^{(y)}(t_2) &= \frac{1}{2} E_0 e^{-i\omega_L t_{0y}} = \\ &= \frac{1}{2} E_0 e^{-i\omega_L(t_2 - 2L) - i\phi_0 + i\Delta\phi_y} \end{aligned} \quad (89)$$

with $\Delta\phi_y = -\Delta\phi_x$. The fields E_{pd} recombined at the photo-detector gives

$$\begin{aligned} E_{pd}(t) &= E^{(x)}(t) + E^{(y)}(t) \\ &= -iE_0 e^{-i\omega_L(t-2L)} \sin(\phi_0 + \Delta\phi_x) , \end{aligned} \quad (90)$$

with a total power $P = P_0 \sin^2(\phi_0 + \Delta\phi_x(t))$. Note that at the other output of the beam splitter, towards the laser, $E_L = E^{(x)} - E^{(y)}$ so that energy is conserved. The optimal length giving the highest $\Delta\phi_x$ is

$$L = \frac{\pi}{2\omega_{GW}} \simeq 750\text{km} \left(\frac{f_{GW}}{100\text{Hz}} \right)^{-1}. \quad (91)$$

Actually real interferometers include Fabry-Perot cavities, where the laser beam goes back and forth several times before being recombined at the beam splitter, allowing the *actual* of the photon travel path to be ~ 100 km. For discussion of real interferometers with Fabry-Perot cavities see e.g. sec. 9.2 of ²², with the result that the sensitivity is enhanced by a factor $2F/\pi$ where F is the finesse of the Fabry-Perot cavity and typically $F \sim O(100)$, giving a measured phase-shift of the order

$$\Delta\phi_{FB} \sim \frac{4F}{\pi} \omega_L L h_0 \simeq \frac{8F}{\pi} \omega_L \Delta L \quad (92)$$

The typical amplitude h_0 that can be measured is of order 10^{-20} which at the best sensitive frequency gives for the Michelson interferometer $\delta L \sim 10^{-15}$ km!

The typical GW amplitude emitted by a binary system is

$$h \sim G_N M v^2 / r \simeq 2.4 \times 10^{-22} \left(\frac{M}{M_\odot} \right) \left(\frac{v}{0.1} \right)^2 \left(\frac{r}{\text{Mpc}} \right)^{-1} \quad (93)$$

many order of magnitudes small than the earth gravitational field. It is its peculiar time oscillating behaviour that makes possible its detection. We will see however in sec. that rather than the instantaneous amplitude of the signal, its integrated value will be of interest for GW detection.

Numerology

Interferometric detectors are very precise and rapidly responsive ruler, they can detect the change of an arm length down to values of 10^{-15} m, however not at all frequency scales. At very low frequency ($f_{GW} \lesssim 10$ Hz) the noise from seismic activity and generic vibrations degrade the sensitivity of the instrument, whereas at high frequency laser shot noise does not allow to detect signal with frequency larger than few kHz. Considering binary systems, which emit according to the flux given in eq. (3), what is the typical length, mass, distance scale of the source? Using eq. (4) we obtain

$$\begin{aligned} v &= (G_N M \pi f_{GW})^{1/3} \simeq 0.054 \left(\frac{M}{M_\odot} \right)^{1/3} \left(\frac{f_{GW}}{10\text{Hz}} \right)^{1/3}, \\ r &= G_N M (G_N M \pi f_{GW})^{-2/3} \simeq 6.4\text{Km} \left(\frac{M}{M_\odot} \right)^{1/3} \left(\frac{f_{GW}}{10\text{Hz}} \right)^{-2/3}. \end{aligned} \quad (94)$$

²² M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008

It can also be interesting to estimate how long it will take for a coalescence to take place. Using the lowest order expression for energy and flux, one has

$$\begin{aligned} -\eta M v \frac{dv}{dt} &= -\frac{32}{5G_N} \eta^2 v^{10} \implies \\ \frac{dv}{v^9} &= \frac{32\eta}{5G_N M} dt \implies \\ \frac{1}{v_i^8} - \frac{1}{v_f^8} &= \frac{256\eta}{5G_N M} \Delta t, \end{aligned} \quad (95)$$

for the time Δt taken for the inspiralling system to move from v_i to v_f . If $v_i \ll v_f$ we can estimate

$$\Delta t \simeq \frac{5G_N M}{256\eta} v_i^{-8} \simeq 1.4 \times 10^4 \text{sec} \frac{1}{\eta} \left(\frac{M}{M_\odot} \right)^{-5/3} \left(\frac{f_{i\text{GW}}}{10\text{Hz}} \right)^{-8/3}. \quad (96)$$

Note that v_f can be comparable to v_i for very massive systems, which enter the detector sensitivity band when $v_i \lesssim 1$. For an estimate of the maximum relative binary velocity during the inspiral, we can take the inner-most stable circular orbit v_{ISCO} of the Schwarzschild case, which gives

$$v_{\text{ISCO}} = \frac{1}{\sqrt{6}} \simeq 0.41. \quad (97)$$

The number of cycles N the GW spends in the detector sensitivity band can be derived by noting that

$$E = -\frac{1}{2} \eta M (G_N M \pi f_{\text{GW}})^{2/3} \quad (98)$$

and from eq. (5)

$$\begin{aligned} N(t) &= \int_{t_i}^t f_{\text{GW}}(t) dt' \implies \\ N(f_{\text{GW}}) &\simeq \int_{f_{i\text{GW}}}^{f_{\text{GW}}} f \frac{dE/df}{dE/dt} df \\ &\simeq \frac{5G_N M}{96\eta} \int_{f_{i\text{GW}}}^{f_{\text{GW}}} (G_N M \pi f)^{-8/3} df \\ &= \frac{1}{32\pi\eta} (G_N M \pi)^{-5/3} \left(\frac{1}{f_{i\text{GW}}^{-5/3}} - \frac{1}{f_{\text{GW}}^{-5/3}} \right) \\ &\simeq 1.5 \times 10^5 \frac{1}{\eta} \left(\frac{M}{M_\odot} \right)^{-5/3} \left(\frac{f_{i\text{GW}}}{10\text{Hz}} \right)^{-5/3} \end{aligned} \quad (99)$$

We can finally obtain the time evolution of the GW frequency

$$\dot{f}_{\text{GW}} = \frac{96}{5} \pi^{8/3} \eta (G_N M)^{5/3} f_{\text{GW}}^{11/3} \quad (100)$$

which has solution

$$\begin{aligned} f_{\text{GW}}(t) &= \frac{1}{\eta^{3/8} \pi} \left(\frac{5}{256} \frac{1}{|t|} \right)^{3/8} G_N M^{-5/8} \\ &= 151 \text{Hz} \frac{1}{\eta^{3/8}} \left(\frac{M}{M_\odot} \right)^{-5/8} \left(\frac{|t|}{1\text{sec}} \right)^{-3/8}, \end{aligned} \quad (101)$$

which can be inverted to give

$$|t_*(f)| = \frac{5}{256\pi\eta} \left(\frac{1}{\pi G_N M \pi} \right)^{5/3}. \quad (102)$$

Exercise 1 ** Linearized Riemann, Ricci and Einstein tensors

Using that the Christoffel symbols at linear level are

$$\Gamma_{\mu\nu}^{\alpha} = \frac{1}{2} \left(\partial_{\mu} h_{\nu}^{\alpha} + \partial_{\nu} h_{\mu}^{\alpha} - \partial^{\alpha} h_{\mu\nu} \right)$$

derive eqs.(12)

Exercise 2 * Retarded Green function I**

Show that the two representation of the retarded Green function given by eq. (17) and

$$G_{ret}(t, x) = -i\theta(t) (\Delta_{+}(t, x) - \Delta_{-}(t, x)) ,$$

where

$$\Delta_{\pm}(t, x) \equiv \int_{\mathbf{k}} e^{\mp ikt} \frac{e^{i\mathbf{k}\mathbf{x}}}{2k}$$

are equivalent. Hint: use that

$$\int_{-\infty}^{\infty} \frac{dk}{2\pi} e^{ikx} = \delta(x) ,$$

and that

$$\theta(t) \int_{-\infty}^{\infty} \frac{dk}{2\pi} e^{ik(t+r)} = 0 \quad \text{for } r \geq 0 .$$

Exercise 3 * Retarded Green function II**

Use the representation of the G_{ret} obtained in the previous exercise to show that

$$G_{ret}(t, \mathbf{x}) = - \int_{\mathbf{k}} \frac{d\omega}{2\pi} \frac{e^{-i\omega t + i\mathbf{k}\mathbf{x}}}{k^2 - (\omega + i\epsilon)^2} ,$$

where ϵ is an arbitrarily small positive quantity. Hint: use that

$$\theta(\pm t) = \mp \frac{1}{2\pi i} \int \frac{e^{-i\omega t}}{\omega \pm i\epsilon} .$$

Show that G_{ret} is real.

Exercise 4 * Advanced Green function**

Same as the two exercises above for G_{adv} , with

$$\begin{aligned} G_{adv}(t, \mathbf{x}) &= i\theta(-t) (\Delta_{+}(t, x) - \Delta_{-}(t, x)) , \\ G_{adv}(t, \mathbf{x}) &= - \int_{\mathbf{k}} \frac{d\omega}{2\pi} \frac{e^{-i\omega t + i\mathbf{k}\mathbf{x}}}{k^2 - (\omega - i\epsilon)^2} . \end{aligned}$$

Show that G_{adv} is real.

Exercise 5 * Feynman Green function I**

Show that the G_F defined by

$$G_F(t, \mathbf{x}) = -i \int_{\mathbf{k}} \frac{d\omega}{2\pi} \frac{e^{-i\omega t + i\mathbf{k}\mathbf{x}}}{k^2 - \omega^2 - i\epsilon}$$

is equivalent to

$$G_F(t, \mathbf{x}) = \theta(t)\Delta_+(t, \mathbf{x}) + \theta(-t)\Delta_-(t, \mathbf{x}).$$

Derive the relationship

$$G_F(t, \mathbf{x}) = \frac{i}{2} (G_{adv}(t, \mathbf{x}) + G_{ret}(t, \mathbf{x})) + \frac{\Delta_+(t, \mathbf{x}) + \Delta_-(t, \mathbf{x})}{2}.$$

Exercise 6 *** Feynman Green function II

By integrating over \mathbf{k} the G_F in the $\sim 1/(k^2 - \omega^2)$ representation, show that G_F implements boundary conditions giving rise to field h behaving as

$$h(t, \mathbf{x}) \sim \int d\omega e^{-i\omega t + i|\omega|r},$$

corresponding to out-going (in-going) wave for $\omega > (<)0$. Since an $\omega < 0$ solution is equivalent to a $\omega > 0$ solution propagating backward in time, this result can be interpreted by saying that using G_F results into having pure out-going (in-going) wave for $t \rightarrow \pm\infty$.

Exercise 7 * TT gauge

Show that the projectors defined in eq. 22 satisfy the relationships

$$\begin{aligned} P_{ij}P_{jk} &= P_{ik} \\ \Lambda_{ij,kl}\Lambda_{kl,mn} &= \Lambda_{ij,mn}, \end{aligned}$$

which characterize projectors operator.

Exercise 8 ***** Energy of circular orbits in a Schwarzschild metric

Consider the Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2G_N M}{r}\right) dt^2 + \frac{dr^2}{\left(1 - \frac{2G_N M}{r}\right)} + r^2 d\Omega^2. \quad (103)$$

The dynamics of a point particle with mass m moving in such a background can be described by the action

$$S = -m \int d\tau = -m \int d\lambda \sqrt{-g_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}}$$

for any coordinate λ parametrizing the particle world-line. Using $S = \int d\lambda L$, we can write

$$L = -m \left[\left(1 - \frac{2G_N M}{r}\right) \left(\frac{dt}{d\tau}\right)^2 - \frac{\left(\frac{dr}{d\tau}\right)^2}{\left(1 - \frac{2G_N M}{r}\right)} - r^2 \left(\frac{d\phi}{d\tau}\right)^2 \right]^{1/2}.$$

Verify that L has cyclic variables t and ϕ and derive the corresponding conserved momenta.

(Hint: use $g_{\mu\nu}(dx^\mu/d\tau)(dx^\nu/d\tau) = -1$. Result: $E = m(dt/d\tau)(1 - 2G_N M/r) \equiv m e$ and $L = mr^2(d\phi/d\tau) \equiv m l$).

By expressing $dt/d\tau$ and $d\phi/d\tau$ in terms of e and l , derive the relationship

$$e^2 = (1 - 2G_N M/r) \left(1 + l^2/r^2\right) + \left(\frac{dr}{d\tau}\right)^2.$$

From the circular orbit conditions ($\frac{de}{dr} = 0 = dr/d\tau = 0$), derive the relationship between l and r for circular orbits.

(Result: $l^2 = r^2 / (\frac{r}{M} - 3)$).

Substitute into the energy function e and find the circular orbit energy

$$e(x) = \frac{1 - 2x}{\sqrt{1 - 3x}},$$

where $x \equiv (G_N M \dot{\phi})^{2/3}$ is an observable quantity as it is related to the GW frequency f_{GW} by $x = (G_N M \pi f_{GW})^{2/3}$.

(Hint: Use

$$\dot{\phi} = \frac{d\phi}{d\tau} \dot{\tau} = \frac{l}{r^2} \left[1 - \frac{2M}{r} - r^2 \dot{\phi}^2\right]^{1/2}$$

to find that on circular orbits $(M\dot{\phi})^2 = (G_N M/r)^3$, an overdot stands for derivative with respect to t .)

Derive the relationships for the Inner-most stable circular orbit

$$\begin{aligned} r_{ISCO} &= 6G_N M = 4.4\text{km} \left(\frac{M}{M_\odot}\right) \\ f_{ISCO} &= \frac{1}{6^{3/2}} \frac{1}{G_N M \pi} \simeq 8.8\text{kHz} \left(\frac{M}{M_\odot}\right)^{-1} \\ v_{ISCO} &= \frac{1}{\sqrt{6}} \simeq 0.41 \end{aligned}$$

Exercise 9 ** Ruler under GW action

Derive the solution (80) to the eq. (79).

Exercise 10 ** Newtonian force exerted by GWs

Derive eq. (83) from eq. (82)

Exercise 11 **** Energy released by GWs

Derive the work done on the experimental apparatus by the GW Newtonian force of eq. (83).

The post-Newtonian expansion in the effective field theory approach

We want to obtain the PN correction to the Newtonian potential due to GR and we want to work at the level of the *equation of motions*. The dynamics for the massive particle (star/black hole) is given by the world-line action

$$\begin{aligned}
 S &= -m \int d\tau \\
 &= -m \int d\tau \sqrt{-g_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau}} \\
 &= -m \int dt \left[-g_{00} - 2g_{0i} \frac{dx^i}{dt} - g_{ij} \frac{dx^i}{dt} \frac{dx^j}{dt} \right]^{1/2}
 \end{aligned} \tag{104}$$

and the dynamics of the gravitational degrees of freedom is given by the standard Einstein-Hilbert action, expanded to quadratic order

$$S_{EH} = -\frac{1}{64\pi G_N} \int dt d\mathbf{x} \left[\partial_\mu h_{\alpha\beta} \partial^\mu h^{\alpha\beta} - \partial_\mu h \partial^\mu h + 2\partial_\mu h^{\mu\nu} \partial_\nu h - 2\partial_\mu h^{\mu\nu} \partial_\rho h_\nu^\rho \right] \tag{105}$$

Working at the level of e.o.m. we could solve them perturbatively, by taking as a first approximation ²³

$$\square \bar{h}_{\mu\nu} = -16\pi G_N T_{\mu\nu}, \tag{106}$$

²³ Note that left hand side of eq. (15) does not follow from eq. (105), as the gauge fixing term (109) is missing.

then using the solution

$$\bar{h}_{\mu\nu}^{(N)} = 4G_N \int dt d^3\mathbf{x}' G_{Ret}(t-t', \mathbf{x}-\mathbf{x}') T_{\mu\nu}(t', \mathbf{x}') \tag{107}$$

and finally pluggin this solution back into the $O(h^2)$ Einstein equation

$$\square h_{\mu\nu}^{(1PN)} \simeq \partial^2 \left(h_{\mu\nu}^{(N)} \right)^2 \tag{108}$$

but we are going to perform the computation more efficiently.

First we have to impose the gauge condition. The action (105) has to be completed with a *gauge fixing* term S_{GF} given by

$$S_{GF} = \frac{1}{32\pi G_N} \int dt d\mathbf{x} \left(\partial_\nu h_{\mu\nu} - \frac{1}{2} \partial_\mu h \right)^2 \tag{109}$$

taking the total action to

$$S_{EH+GFF} = \frac{1}{64\pi G_N} \int dt d^d x h^{\mu\nu} A_{\mu\nu\rho\sigma} \square h^{\rho\sigma} \quad (110)$$

with

$$A_{\mu\nu\rho\sigma} = \frac{1}{2} (\eta_{\mu\rho}\eta_{\nu\sigma} + \eta_{\mu\sigma}\eta_{\nu\rho} - \eta_{\mu\nu}\eta_{\rho\sigma}) . \quad (111)$$

In order to find the Green function for h we need to invert the A operator, which is straightforwardly done by noting that

$$A_{\mu\nu\alpha\beta} A^{\alpha\beta}_{\rho\sigma} = \frac{1}{2} (\eta_{\mu\rho}\eta_{\nu\sigma} + \eta_{\mu\sigma}\eta_{\nu\rho}) . \quad (112)$$

We can use this Lagrangian construction in order to solve for the h field, as in ex. 14, but here we want to show how powerful the effective action method is in determining the dynamics of the 2-body system, “integrating out” the gravitational degrees of freedom.

Let us pause briefly to introduce some technicalities about *Gaussian integrals* that will be helpful later. The basic formula we will need is

$$\int_{-\infty}^{\infty} e^{-\frac{1}{2}ax^2+Jx} dx = \left(\frac{2\pi}{a}\right)^{1/2} \exp\left(\frac{J^2}{2a}\right) . \quad (113)$$

or its multi-dimensional generalization

$$\int e^{-\frac{1}{2}x^i A_{ij} x^j + J^i x_i} dx_1 \dots dx_n = \frac{(2\pi)^{n/2}}{(\det A)^{1/2}} \exp\left(\frac{1}{2} J^t A^{-1} J\right) . \quad (114)$$

Other useful formulae are

$$\int x^k e^{-\frac{1}{2}ax^2+Jx} dx = \left(\frac{2\pi}{a}\right)^{1/2} \left(\frac{d}{dJ}\right)^k \exp\left(\frac{J^2}{2a}\right) \quad (115)$$

from which it follows that

$$\begin{aligned} \int x^{2n} e^{-\frac{1}{2}ax^2} dx &= \left(\frac{2\pi}{a}\right)^{1/2} \left(\frac{d}{dJ}\right)^{2n} \exp\left(\frac{J^2}{2a}\right) \Big|_{J=0} \\ &= \frac{(2n-1)!!}{a^n} \left(\frac{2\pi}{a}\right)^{1/2} , \end{aligned} \quad (116)$$

which also admit a natural generalization in case of x is not a real number but an element of a vector space. Let us see how this will be useful in the toy model of massless, non self-interacting scalar field Φ interacting with a source J :

$$\begin{aligned} S_{toy} &= \int dt d^d x \left[-\frac{1}{2} (\partial\Phi(t, \mathbf{x}))^2 + J(t, \mathbf{x})\Phi(t, \mathbf{x}) \right] \\ &= \int_{\mathbf{k}} \frac{dk_0}{2\pi} \left[\Phi(k_0, \mathbf{k})\Phi^*(k_0, \mathbf{k}) (k_0^2 - k^2) + J(k_0, \mathbf{k})\Phi^*(k_0, \mathbf{k}) \right] \end{aligned} \quad (117)$$

and apply the above eqs.(113–116), with two differences:

- Here the integration variable is Φ , depending on 2 continuous indices (k_0, \mathbf{k}) , instead of the discrete index $i \in 1 \dots n$
- the Gaussian integrand is actually turned into a complex one, as we are taking at the exponent

$$Z_0[J] \equiv \int \mathcal{D}\Phi \exp \left\{ i \int_{\mathbf{k}} \frac{dk_0}{2\pi} \left[\frac{1}{2} (k_0^2 - k^2) \Phi(k_0, \mathbf{k}) \Phi(-k_0, -\mathbf{k}) + J(k_0, \mathbf{k}) \Phi(-k_0, -\mathbf{k}) + i\epsilon |\Phi(k_0, \mathbf{k})|^2 \right] \right\}, \quad (118)$$

where the ϵ has been added term ensure convergence for $|\Phi| \rightarrow \infty$.

We can now perform the Gaussian integral by using the new variable

$$\Phi'(k_0, \mathbf{k}) = \Phi(k_0, \mathbf{k}) + (k_0^2 - k^2 + i\epsilon) J(k_0, \mathbf{k})$$

that allows to rewrite eq. (118) as

$$Z_0[J] = \exp \left[-\frac{i}{2} \int_{\mathbf{k}} \frac{dk_0}{2\pi} \frac{J(k_0, \mathbf{k}) J^*(k_0, \mathbf{k})}{k_0^2 - k^2 + i\epsilon} \right] \times \int \mathcal{D}\Phi' \exp \left\{ i \int_{\mathbf{k}} \frac{dk_0}{2\pi} \left[\frac{1}{2} (k_0^2 - k^2) \Phi'(k_0, \mathbf{k}) \Phi'^*(k_0, \mathbf{k}) + i\epsilon |\Phi'|^2 \right] \right\}. \quad (119)$$

The integral over Φ' gives an uninteresting normalization factor \mathcal{N} , thus we can write the result of the functional integration as

$$\begin{aligned} Z_0[J] &= \mathcal{N} \exp \left[-\frac{i}{2} \int_{\mathbf{k}} \frac{dk_0}{2\pi} \frac{J(k_0, \mathbf{k}) J(-k_0, -\mathbf{k})}{k_0^2 - k^2 + i\epsilon} \right] \\ &= \mathcal{N} \exp \left[-\frac{1}{2} \int dt d^3x G_F(t-t', \mathbf{x}-\mathbf{x}') J(t, \mathbf{x}) J(t', \mathbf{x}') \right], \end{aligned} \quad (120)$$

The $Z_0[J]$ functional is the main ingredient allowing to compute the *effective action* describing the dynamics of the sources of the field we are integrating over and the dynamics of the extra field we are *not* integrating over. For instance starting from S_{toy} defined in eq. (117) we would obtain the effective action for the sources J from

$$S_{eff}[J] = -i \log Z_0[J], \quad (121)$$

where we can safely discard the normalization constant \mathcal{N} . For instance substituting in S_{toy} $J\phi \rightarrow J_0 + J\Phi$, with

$$J_0(t, \mathbf{x}) + J(t, \mathbf{x}) \Phi(t, \mathbf{x}) = - \sum_A m_A \delta(t - \tau_A) \delta^{(3)}(\mathbf{x} - \mathbf{x}_A) (1 + \Phi(t, \mathbf{x})) \quad (122)$$

one would obtain the effective action

$$\begin{aligned} S_{eff}(x_A) &= \sum_A \left[-m_A \int d\tau_A + \right. \\ &\quad \left. \frac{i}{2} \sum_B m_A m_B \int d\tau_A d\tau_B G_F(t(\tau_A) - t(\tau_B), \mathbf{x}_A(\tau_A) - \mathbf{x}_B(\tau_B)) \right]. \end{aligned} \quad (123)$$

Taking for the Green function the quasi static limit

$$\begin{aligned}
& -i \int_{\mathbf{k}} \frac{d\omega}{2\pi} \frac{e^{-i\omega(t_A-t_B)+i\mathbf{k}(\mathbf{x}_A-\mathbf{x}_B)}}{k^2 - \omega^2 + i\epsilon} \\
& \simeq -i \int_{\mathbf{k}} \frac{d\omega}{2\pi} \frac{e^{-i\omega(t_A-t_B)+i\mathbf{k}(\mathbf{x}_A-\mathbf{x}_B)}}{k^2} \left(1 + \frac{\omega^2}{k^2} + \dots \right) \\
& \simeq -i\delta(t_A - t_B) \int_{\mathbf{k}} \frac{e^{i\mathbf{k}(\mathbf{x}_A-\mathbf{x}_B)}}{\mathbf{k}^2} \left(1 + \frac{\partial_{t_1}\partial_{t_2}}{\mathbf{k}^2} \right) \\
& = -i\delta(t) \left[\frac{1}{4\pi|\mathbf{x}|} + \frac{O(v^2)}{|\mathbf{x}|} \right]
\end{aligned} \tag{124}$$

one recover the instantaneous $1/r$, Newtonian interaction (plus $O(v^2)$ corrections). Note that we have implemented the substitution $\omega = -i\partial_{t_1} = i\partial_{t_2}$ in order to work out the systematic expansion in v . This is justified by observing that the wave-number $k^\mu \equiv (k^0, \mathbf{k})$ of the gravitational modes mediating this interaction have ($k^0 \sim v/r, k \sim 1/r$), so in order to have manifest power counting it is necessary to Taylor expand the propagator.

The individual particles can also exchange *radiative* gravitons (with $k_0 \simeq k \sim v/r$), but such processes give sub-leading contributions to the effective potential in the PN expansion, and they will be dealt with in sec. . In other words we are not integrating out the entire gravity field, but the specific off-shell modes in the kinematic region $k_0 \ll k$.

Actually there are some more terms we would have obtained, like the $J_{A,B}^2 G_F(0,0)$ which are divergent, as they involve the Green function computed at 0 separation in space-time. These corresponds to a source interacting with itself and we can safely discard it, as such term does not contribute in any way to the 2-body potential. Its unobservable (infinite) contribution can be re-absorbed by an (infinite) shift of the value of the mass (any ultraviolet divergence can be reabsorbed by a *local* counter term). Of course we cannot trust our theory at arbitrarily short distance, where this divergence may be regularized by new physics (quantum gravity?) but as we do not aim to *predict* the parameters of the theory, but rather take them as input to compute other quantities like interaction potential, we keep intact the predictive power of our approach. From the technical point of view it is a power-law divergence, which in *dimensional regularization* is automatically set to zero.

If there are interaction terms which cannot be written with terms linear or quadratic in the field, the Gaussian integral cannot be done analytically, so the rule to follow is to separate the quadratic action $S_{quad}[\Phi]$ of the field (its kinetic term) and Taylor expand all the rest:

for an action $S = S_{quad} + (S - S_{quad})$ one would write

$$\begin{aligned} Z[J] &= \int \mathcal{D}\Phi e^{iS+i \int J\Phi} \\ &= \int \mathcal{D}\Phi e^{iS_{quad}+i \int J\Phi} \left[1 + i(S - S_{quad}) - \frac{(S - S_{quad})^2}{2} + \dots \right], \end{aligned} \quad (125)$$

where J is now an auxiliary source, the physical source term will be Taylor expanded in the $S - S_{quad}$ term. As long as the $S - S_{quad}$ contains only polynomials of the field which is integrated over, the integral can be *perturbatively* performed analytically, inheriting the rule from eqs. (115,116): roughly speaking fields have to be paired up, each pair is going to be substituted by a Green function.

Our perturbative expansion admits a nice and powerful representation in terms of Feynman diagrams. Incoming and outgoing particle world-lines are represented by horizontal lines, Green functions by dashed lines connecting points, see e.g. fig. 2 the Feynman diagram accounting for the Newtonian potential, which is obtained by pairing the fields connected in the following expression

$$\begin{aligned} e^{iS_{eff}} &= Z[J, x_A]|_{J=0} = \int \mathcal{D}\Phi e^{iS_{quad}} \times \left\{ 1 \right. \\ &\quad \left. - \frac{1}{2} \left[\sum_A m_A \int dt_A \Phi(t_A, \mathbf{x}_A(t_A)) \right] \left[\sum_B m_B \int dt_B \Phi(t_B, \mathbf{x}_B(t_B)) \right] + \dots \right\} \end{aligned} \quad (126)$$




Figure 2: Feynman graph accounting for the Newtonian potential.

If following Green function's lines all the vertices can be connected the diagram is said *connected*, otherwise it is said *disconnected*: only connected diagrams contribute to the effective action. We will not demonstrate this last statement, but its proof relies on the following argument. Taking the logarithm of eq. (126) we get

$$S_{eff} = -i \log Z_0[0] - i \log \left(\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} Z_0^{-1}[0] (Z[0] - Z_0[0])^n \right). \quad (127)$$

All terms with $n > 1$ describe disconnected diagrams, and some disconnected diagrams can also be present in the $n = 1$ term. However the $n = 1$ disconnected contribution is precisely canceled by the

$n = 2$ terms. Beside discarding the disconnected diagrams, we can also discard diagrams involving Green function at vanishing separation $G_F(0,0)$, which give an infinite re-normalization to parameters with dimensions. For instance the diagram in fig. 3, which would arise from the $(S_{quad} - S)^4$ term in eq. (126), is both disconnected and involve a $G_F(0,0)$, cancelling the contribution from the product of disconnected diagrams from $(Z - Z_0)^2$, where in one factor of $Z - Z_0$ one contracts two ϕ s at equal point, and in the other two ϕ s at different points.




Figure 3: Diagram giving a power-law divergent contribution to the mass.

Having exposed the general method, let us apply it to the computation of the effective action for the conservative dynamics of binary systems. In order to do that it will be useful to decompose the metric as

$$g_{\mu\nu} = e^{2\phi/\Lambda} \begin{pmatrix} -1 & A_j/\Lambda \\ A_i/\Lambda & e^{-c_d\phi/\Lambda}\gamma_{ij} - A_iA_j/\Lambda^2 \end{pmatrix}, \quad (128)$$

with $\gamma_{ij} = \delta_{ij} + \sigma_{ij}/\Lambda$, $c_d = 2\frac{(d-1)}{(d-2)}$ ($\Lambda = (32\pi G_N)^{-1/2}$ is a constant with dimensions that will allow a simpler normalization of the Green functions). In terms of this parametrization, the Einstein-Hilbert plus gauge fixing action is at quadratic order

$$S_{quad} = \int dt d^d x \sqrt{\gamma} \left\{ \frac{1}{4} \left[(\vec{\nabla}\sigma)^2 - 2(\vec{\nabla}\sigma_{ij})^2 - (\dot{\sigma}^2 - 2(\dot{\sigma}_{ij})^2) \right] - c_d \left[(\vec{\nabla}\phi)^2 - \dot{\phi}^2 \right] + \left[\frac{F_{ij}^2}{2} + (\vec{\nabla}\cdot\vec{A})^2 - \dot{A}^2 \right] \right\}, \quad (129)$$

where the gauge fixing term

$$S_{GF} = \frac{1}{32\pi G_N} \int dt d^d x \left(g_{il} \tilde{\Gamma}_{jk}^i \tilde{\Gamma}_{mn}^l g^{jk} g^{mn} \right) \quad (130)$$

with $\tilde{\Gamma}_{jk}^i \equiv \frac{1}{2}\gamma^{il} (\gamma_{ljk} + \gamma_{lkj} - \gamma_{jk,l})$ has been used, and the source

term is

$$\begin{aligned}
S_p &= -m \int dt e^{\phi/\Lambda} \sqrt{\left(1 - \frac{A_i v^i}{\Lambda}\right) - e^{-c_d \phi/\Lambda} (v^2 + \sigma_{ij} v^i v^j)} \\
&\simeq -m \int dt \left\{ \sqrt{1 - v^2} + \phi \left[1 + \frac{3}{2} v^2\right] + \frac{\phi^2}{2\Lambda} \left[1 + O(v^2)\right] + \right. \\
&\quad \left. + A_i v_i \left[1 + O(v^2)\right] + \frac{\sigma_{ij}}{2} v_i v_j \left[1 + O(v^2)\right] + \dots \right\}. \quad (131)
\end{aligned}$$

The Green functions (inverse of the quadratic terms) of the fields are

$$G_F(t, \mathbf{x}) = -i \int_{\mathbf{k}} \frac{dk_0}{2\pi} \frac{e^{-ik_0 t + i\mathbf{k}\mathbf{x}}}{k^2 - k_0^2} \times \begin{cases} -\frac{1}{2c_d} & \phi \\ \frac{1}{2} \delta_{ij} & A \\ -\frac{1}{2} \left(\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk} - \frac{2}{d-2} \delta_{ij} \delta_{kl} \right) & \sigma \end{cases} \quad (132)$$

Let us consider the computation of the effective action

$$\begin{aligned}
iS_{eff} &= \log \int \mathcal{D}\phi \mathcal{D}A \mathcal{D}\sigma e^{iS_{quad}} \left\{ 1 + \dots - \frac{1}{2} \right. \\
&\quad \times \left[-\frac{m_1}{\Lambda} \int dt_1 \left(\phi \left(1 + \frac{3}{2} v_1^2\right) + \frac{\phi^2}{2\Lambda} + A_i v_{1i} \right) \right. \\
&\quad \left. \left. - \frac{m_2}{\Lambda} \int dt_2 \left(\phi \left(1 + \frac{3}{2} v_2^2\right) + \frac{\phi^2}{2\Lambda} + A_i v_{2i} \right) + \dots \right]^2 \right. \\
&\quad \left. + \frac{i^3}{6} \left[-\frac{m_1}{\Lambda} \int dt_1 (\phi(1 + \dots)) - \frac{m_2}{\Lambda} \int dt_2 \left(\dots + \frac{\phi^2}{2\Lambda} + \dots \right) \right]^3 + \dots \right\}, \quad (133)
\end{aligned}$$

We see that we have to pair up, or *contract* the term linear in ϕ of the second line with the analog term in the third line, to give a contribution to the effective action

$$\begin{aligned}
iS_{eff}|_{fig. 2-\phi} &\supset -i^3 \frac{m_1 m_2}{8\Lambda^2} \int dt_1 dt_2 \delta(t_1 - t_2) \int_{\mathbf{k}} \frac{e^{i\mathbf{k}(\mathbf{x}_1(t_1) - \mathbf{x}_2(t_2))}}{k^2} \left[1 + \frac{3}{2} (v_1^2 + v_2^2)\right] \left(1 + \frac{\partial_{t_1} \partial_{t_2}}{k^2}\right) \\
&= i \frac{m_1 m_2}{8\Lambda^2} \int dt \int_{\mathbf{k}} \frac{e^{i\mathbf{k}(\mathbf{x}_1(t) - \mathbf{x}_2(t))}}{k^2} \left[1 + \frac{3}{2} (v_1^2 + v_2^2)\right] \left(1 + \frac{v_1^i v_2^j k_i k_j}{k^2}\right) \\
&\simeq i \frac{G_N m_1 m_2}{r} \left[1 + \frac{3}{2} (v_1^2 + v_2^2) + \frac{1}{2} (v_1 v_2 - (v_1 \hat{r})(v_2 \hat{r}))\right] \quad (134)
\end{aligned}$$

where the propagator has been Taylor expanded around $k_0/k \sim 0$ as in eq. (124) and the formulae

$$\int_{\mathbf{k}} e^{i\mathbf{k}\mathbf{x}} \frac{1}{k^{2\alpha}} = \frac{\Gamma(d/2 - \alpha)}{(4\pi)^{d/2} \Gamma(\alpha)} \left(\frac{r}{2}\right)^{2\alpha - d} \quad (135)$$

$$\int_{\mathbf{k}} e^{i\mathbf{k}\mathbf{x}} \frac{k^i k^j}{k^{2\alpha}} = \left(\frac{1}{2} \delta^{ij} - \left(\frac{d}{2} - \alpha + 1\right) \hat{r}^i \hat{r}^j\right) \frac{\Gamma(d/2 - \alpha + 1)}{(4\pi)^{d/2} \Gamma(\alpha)} \left(\frac{r}{2}\right)^{2\alpha - d - 2} \quad (136)$$

have been used. Considering the contraction of two A fields one gets

$$\begin{aligned}
iS_{eff}|_{fig. 2-A_i} &\supset i^3 \frac{m_1 m_2}{2\Lambda^2} \int dt \int_{\mathbf{k}} \frac{e^{i\mathbf{k}(\mathbf{x}_1 - \mathbf{x}_2)}}{k^2} \delta_{ij} v_1^i v_2^j \\
&= -i \frac{4G_N m_1 m_2}{r} v_1 v_2 \quad (137)
\end{aligned}$$

This is still not the whole story for the v^2 corrections to the Newtonian potential, as we still have to contract two ϕ 's from the fourth line of eq. (133) with a ϕ^2 of the same line, getting the contribution

$$\begin{aligned} iS_{eff}|_{fig. 4} &\supset i^5 \frac{m_1^2 m_2}{128\Lambda^4} \int dt \left(\int_{\mathbf{k}} \frac{e^{i\mathbf{k}(\mathbf{x}_1 - \mathbf{x}_2)}}{k^2} \right)^2 \\ &= i \frac{G_N^2 m_1^2 m_2}{2r^2} \end{aligned} \quad (138)$$

Summing the contributions from eqs.(134,137,138), plus the 1 \leftrightarrow 2 of eq. (138), one obtains the Einstein-Infeld-Hoffman potential (remember that the potential enters the Lagrangian with a minus sign!)

$$V_{EIH} = -\frac{G_N m_1 m_2}{2r} \left[3 \left(v_1^2 + v_2^2 \right) - 7v_1 v_2 - (v_1 \hat{r})(v_2 \hat{r}) \right] + \frac{G_N^2 m_1 m_2 (m_1 + m_2)}{2r^2} \quad (139)$$




Figure 4: Graph giving a G_N^2 contribution to the 1PN potential via ϕ propagators.

Power counting

We have nevertheless neglected a contribution from the pairing of the two ϕ^2 terms appearing at $(S_{quad} - S)^2$ order expansion in eq. (133), which gives rise to a term proportional to $G_N m_1 m_2 G_F^2(\mathbf{x}_1 - \mathbf{x}_2)$. We have rightfully discarded it as it represents a *quantum* contribution to the potential, and in the phenomenological situation we are considering to apply this theory, quantum corrections are suppressed with respect to classical terms by terms of the order \hbar/L , where L is the typical angular momentum of the systems, which in our case is

$$L \sim mvr \sim 10^{77} \hbar \left(\frac{m}{M_\odot} \right)^2 \left(\frac{v}{0.1} \right)^{-1}. \quad (140)$$

Intermediate massive object lines, (like the ones in fig. 4) have no propagator associated, as they represent a static source (or sink) of gravitational modes. At the graviton-massive object vertex momentum is *not* conserved, as the graviton momentum is ultra-soft compared to the massive source.

The \hbar counting of the diagrams can be obtained by restoring the proper normalization in the functional action definition eq. (125),

implying that in the expansion we have $[(S - S_{quad})/\hbar]^n$ and that each Green function, being the inverse of the quadratic operator acting on the fields, brings a \hbar^{-1} factor. Note that we consider that the classical sources do not recoil when interacting with via “field pairing”. This is indeed consistent with neglecting quantum effects, as the wavenumber \mathbf{k} of the exchanged gravitational mode has $k \sim 1/r$ and thus momentum \hbar/r .

By inspecting Feynman diagrams we can systematically infer the scaling of their numerical result according to the following rule:

- associate to each n -graviton-particle vertex a factor $m/\Lambda^n dt (d^d k)^n \sim dt m/\Lambda^n r^{-dn}$, and analogously for multiple graviton vertices
- each propagator scales as $\delta(t)\delta^d(k)/k^2 \sim \delta(t)r^{-2+d}$
- each n -graviton internal vertex scale as $(k^2, k_0 k, k_0^2) dt \delta^d(k) (d^d k)^n \sim dt (1, v, v^2) r^{-d(n-1)}$



Figure 5: Vertex scaling: $\frac{m}{\Lambda} dt d^d k \sim dt \frac{m}{\Lambda} r^{-d}$



Figure 6: A Green function is represented by a propagator, with scaling: $\delta(t)\delta^d(k)/k^2 \sim \delta(t)r^{2+d}$



Figure 7: Triple internal vertex scaling: $\frac{(k^2, k k_0, k_0^2)}{\Lambda} \delta^d(k) dt (d^d k)^3 \sim dt \frac{(1, v, v^2)}{r^{2+2d} \Lambda}$

Putting together the previous rule one find for instance that the diagram in fig. 9 scale as dtm (times the appropriate powers of v from the expansion of the vertex and of the propagator)




Figure 8: Quantum contribution to the 2-body potential.

Note that instead of computing the effective potential we can also compute the effective energy momentum tensor of an isolated source by considering the effective action with one external gravitational mode $H_{\mu\nu}(t, x)$ not to be integrated over:

$$\begin{aligned} iS_{eff-1g}(m, H_{\mu\nu}) &= \log \int \mathcal{D}\phi \mathcal{D}A \mathcal{D}\sigma e^{iS_{EH+GF}(h_{\mu\nu}+H_{\mu\nu})} \\ &\quad \times \left(1 - m \int d\tau (h_{\mu\nu} + H_{\mu\nu}) + \dots \right) \Big|_{H_{\mu\nu}^1} \quad (141) \\ &= \frac{i}{2} \int dt d^3x T_{\mu\nu}^{(eff)} H_{\mu\nu}, \end{aligned}$$

where the computation is made as usual by performing a Gaussian integration on all gravity field variables and the result will be linear in the external field which instead of being integrated over, is the $H_{\mu\nu}$ is the one we want to find what it is coupled to. By Lorentz invariance $H_{\mu\nu}$ must be coupled to a symmetric 2-tensor by which by definition is the energy momentum tensor. Alternatively one can take eq. (126) in the presence of physical sources $J \sim -m \int d\tau$ and compute perturbatively the Feynman integral to obtain

$$\langle H_{\mu\nu} \rangle = \int \mathcal{D}\phi \mathcal{D}A \mathcal{D}\sigma H_{\mu\nu} e^{iS(h+H)}. \quad (142)$$

For instance at lowest order it will give

$$\begin{aligned} \langle H_{\mu\nu}(t, x) \rangle &= \int \mathcal{D}\phi \mathcal{D}A \mathcal{D}\sigma H_{\mu\nu} e^{iS_{quad} - im \int dt \frac{\phi}{\Lambda} + H_{00} \dots} \\ &\simeq -i \frac{m}{\Lambda^2} \int dt' d^3y G_F(t-t', x-y) \delta^{(3)}(y-x_1). \quad (143) \end{aligned}$$

By stripping this result by the Green function will give the energy momentum tensor which is coupled to the gravity field, as the solution of Performing this computation at higher perturbative orders will give the higher order corrections to the Newtonian potential.

Exercise 12 *** Geodesic Equation

Derive the geodesic equation from the world line action (104).

Exercise 13 *** Gauge fixed quadratic action

Derive eq. (110) from eq. (105) and the gauge fixing term (109).

Exercise 14 *** Schwarzschild solution at Newtonian order

Starting from action (110) derive the eq. (15) (or one can start with (105 and dropping all terms which *quadratically* vanish on for the Lorentz gauge condition). Expand (104) so to obtain

$$S_{m\text{-static}} = -m \int dt' \left(-1 + \frac{h_{00}(t', \mathbf{x}_p)}{2} \right)$$

and then derive

$$T_{\mu\nu}^{(static)}(t, \mathbf{x}) = 2 \frac{\delta S_{static}}{\delta g^{\mu\nu}(t, \mathbf{x})} = -m \delta^{(3)}(\mathbf{x} - \mathbf{x}_p) \delta_{\mu 0} \delta_{\nu 0}$$

Use eq. (112) to obtain from the variation of the action eq. (15) and plug in the specific form of the retarded Green function to obtain

$$\begin{aligned} \bar{h}_{\mu\nu} &= (-32\pi G_N) \left(-\frac{1}{4\pi} \right) \frac{\delta_{\mu 0} \delta_{\nu 0}}{2} m \int dt' d\mathbf{x}' \frac{\delta(t - t' - |\mathbf{x} - \mathbf{x}'|)}{|\mathbf{x} - \mathbf{x}'|} \delta^{(3)}(\mathbf{x}' - \mathbf{x}_p) \\ &= \frac{4G_N m}{r} \delta_{\mu 0} \delta_{\nu 0}. \end{aligned}$$

Using that

$$h_{\mu\nu} = \bar{h}_{\mu\nu} - \frac{1}{2} \bar{h} \delta_{\mu\nu}$$

and that on the above solution

$$\bar{h} = -4 \frac{G_N m}{r},$$

find the final result

$$\begin{aligned} h_{00} &= \bar{h}_{00} + \frac{1}{2} \bar{h} = 2 \frac{G_N m}{r} \\ h_{xx} &= \bar{h}_{xx} - \frac{1}{2} \bar{h} = 2 \frac{G_N m}{r}. \end{aligned}$$

Exercise 15 * Gaussian integrals

Derive eq. (113) from

$$\int e^{-\frac{1}{2}ax^2} dx = \left(\frac{2\pi}{a} \right)^{1/2}.$$

Derive the above from

$$\left[\int dx e^{-\frac{1}{2}ax^2} \right]^2 = 2\pi \int \rho e^{-\frac{1}{2}a\rho^2} d\rho.$$

Exercise 16 ***** ϕ^3 vertex contribution to the 2PN potential

Compute the contribution of fig. 9 to S_{eff} . The ϕ^3 term in the Einstein-Hilbert Lagrangian is $-c_d \phi \dot{\phi}^2 / \Lambda$, see ²⁴. Hint: this diagram is originated from the $(S - S_{quad})^4$ term which is understood in (133).

Exercise 17 ** Power \hbar

²⁴ Stefano Foffa and Riccardo Sturani. Effective field theory calculation of conservative binary dynamics at third post-Newtonian order. *Phys.Rev.*, D84: 044031, 2011




Figure 9: Sub-leading correction to the 2 particle scattering process due to gravity self-interaction. The ϕ^3 vertex brings two time derivative, making this diagram contribute from 2PN order on.

Derive the \hbar scaling of graphs: \hbar^{I-V} , where I stands for the number of internal lines (propagators) and V for the number of vertices. All classical diagrams are homogeneous in \hbar^{-1}

Exercise 18 *** Quantum correction to classical potential**

Derive the quantum correction to the classical potential given by the term proportional to $G_N m_1 m_2 G_F^2(\mathbf{x}_1 - \mathbf{x}_2)$ originated from the expansion of $S_{quad} - S$ at linear order.

Exercise 19 *** Newtonian potential as graviton exchange**

In non-relativistic quantum mechanics a one-particle state with momentum $\hbar\mathbf{p}$ in the coordinate representation is given by a plane wave

$$\psi_{\mathbf{p}}(\mathbf{x}) = C \exp(i\mathbf{p}\mathbf{x}),$$

where the normalization constant C can be found by imposing

$$\int_V |\psi_{\mathbf{p}}|^2 = 1 \implies C = \frac{1}{\sqrt{V}}.$$

Since we want to trace the powers of M and \hbar in the amplitude, it is necessary to avoid ambiguities: variables \mathbf{p}_i are wave-numbers and ω is a frequency, so that $\hbar\omega$ is an energy and $\hbar\mathbf{p}$ is a momentum.

We define the non-relativistic scalar product

$$\langle \mathbf{p}_1 | \mathbf{p}_2 \rangle^{\text{NR}} = \int d^3x \psi_{\mathbf{p}_1}^*(\mathbf{x}) \psi_{\mathbf{p}_2}(\mathbf{x}) = \delta_{\mathbf{p}_1, \mathbf{p}_2}$$

which differs from the relativistic normalization used in quantum field theory for the scalar product $\langle \mathbf{p}_1 | \mathbf{p}_2 \rangle^{\text{R}}$ according to:

$$\langle \mathbf{p}_1 | \mathbf{p}_2 \rangle^{\text{R}} = \frac{2\omega_{\mathbf{p}_1}}{\hbar} V \langle \mathbf{p}_1 | \mathbf{p}_2 \rangle^{\text{NR}},$$

which implies

$$|\mathbf{p}\rangle^{\text{NR}} = \left(\frac{\hbar}{2\omega_{\mathbf{p}} V} \right)^{1/2} |\mathbf{p}\rangle^{\text{R}}.$$

Note that a scalar field in the relativistic normalization has the *relativistic* expansion in terms of creator and annihilator operators

$$\psi(t, \mathbf{x}) = \int_{\mathbf{k}} \frac{1}{\sqrt{\omega_{\mathbf{k}}}} \left[a_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}t + i\mathbf{k}\mathbf{x}} + a_{\mathbf{k}}^{\dagger} e^{i\omega_{\mathbf{k}}t - i\mathbf{k}\mathbf{x}} \right] \quad (144)$$

with

$$[a_{\mathbf{k}}, a_{\mathbf{k}'}] = i\hbar(2\pi)^3 \delta^{(3)}(\mathbf{k} - \mathbf{k}').$$

We want to study the relativistic analog of the process described in fig. 2, assuming the coupling of gravity with a scalar particle given by the second quantized action

$$S_{pp-quant} = \frac{1}{2} \int dt d^3x \left\{ \partial_{\mu}\psi\partial^{\mu}\psi - \frac{1}{2}\eta_{\mu\nu} \left[(\partial_{\alpha}\psi\partial^{\alpha}\psi) + \frac{M^2}{\hbar^2} \right] \right\} \frac{\hbar^{\mu\nu}}{\Lambda},$$

where the Λ factor has been introduced to have a canonically normalized gravity field.

Derive the quantum amplitude $A(p_1, p_2, p_1 + k, p_2 - k)$ for the 1-graviton exchange between two particles with incoming momenta $p_{1,2}$ and outgoing momenta $p_{1,2} \pm k$

$$\begin{aligned} iA(p_1, p_2, p_1 + k, p_2 - k) &= A_{\alpha\beta\mu\nu} \frac{i\hbar}{k^2 - k_0} \\ &\times \frac{i^2}{\hbar^2 \Lambda^2} \langle \mathbf{p}_1 + \mathbf{k} | T_{\mu\nu} | \mathbf{p}_1 \rangle^{NR} \langle \mathbf{p}_2 - \mathbf{k} | T_{\alpha\beta} | \mathbf{p}_2 \rangle^{NR} \end{aligned}$$

where $A_{\alpha\beta\mu\nu}$ is defined in eq.(111).

Hint: from $S_{p-quant}$ above derive

$$\begin{aligned} T_{00} &= \frac{1}{2} \left[\dot{\psi}^2 + (\partial_i \psi)^2 + \frac{M^2}{\hbar^2} \psi^2 \right], \\ T_{0i} &= \partial_0 \psi \partial_i \psi, \\ T_{ij} &= \partial_i \psi \partial_j \psi - \frac{1}{2} \delta_{ij} \left[-\dot{\psi}^2 + \partial_i \psi \partial^i \psi + \frac{M^2}{\hbar^2} \psi^2 \right], \end{aligned}$$

show that in the non-relativistic limit the term the A_{0000} term dominates and use the above equation to express $|\mathbf{q}\rangle^R$ in terms of $|\mathbf{q}\rangle^{NR}$.

Use also that in the NR limit $p^{\mu} = \delta^{\mu 0} \omega_p = \delta^{\mu 0} M/\hbar$.

Exercise 20 ***** Classical corrections to the Newtonian potential in a quantum set-up

Consider the process in fig. 9, write down its amplitude

$$\begin{aligned} iA_{fig.9} &= \int dt \left(\frac{i}{\hbar} \right)^3 \langle \mathbf{p}_1 + \mathbf{k} | T_{00} | \mathbf{p}_1 \rangle^{NR} \langle \mathbf{p}_2 + \mathbf{q} | T_{00} | \mathbf{p}_2 \rangle^{NR} \langle \mathbf{p}_2 - \mathbf{k} | T_{00} | \mathbf{p}_2 \rangle^{NR} \\ &\times \int_{\mathbf{k}, \mathbf{q}} \frac{d\omega_2}{2\pi} e^{i\mathbf{k}(x_1 - x_2)} \frac{i\hbar}{k^2} \frac{i\hbar}{(\mathbf{k} + \mathbf{q})^2} \frac{i\hbar}{q^2} \frac{i\hbar}{(\mathbf{p}_2 + \mathbf{q})^2 + M^2/\hbar^2 - \omega_2^2 - i\epsilon}. \end{aligned}$$

After writing the massive propagator as

$$\frac{1}{\sqrt{(\mathbf{p}_2 + \mathbf{q})^2 + M^2/\hbar^2 + \omega_2}} \frac{1}{\sqrt{(\mathbf{p}_2 + \mathbf{q})^2 + M^2/\hbar^2 - \omega_2}} \sim \frac{\hbar/M}{\sqrt{(\mathbf{p}_2 + \mathbf{q})^2 + M^2/\hbar^2 - \omega_2}}$$

Perform the integral first in ω_2 and take the limit $\hbar \rightarrow 0$ and $M \rightarrow \infty$ to recover the result in the non-relativistic classical theory.

Exercise 21 ** Correction to Newtonian equation of motions**

Derive the Newtonian-like e.o.m. from the effective action

$$S_{1PN} = \int dt \frac{1}{4} \left(m_1 v_1^4 + m_2 v_2^4 \right) - V_{EIH}.$$

Verify that a result of the type

$$a_1^i = -\omega^2(\mathbf{r}, \mathbf{v})(\mathbf{x}_1^i - \mathbf{x}_2^i) + A(\mathbf{r}, \mathbf{v})(v_1^i - v_2^i)$$

is obtained, for appropriate functions ω and A . Take the circular orbit limit ($\mathbf{r} \cdot \mathbf{v}_{1,2} = 0$) and substitute $x \equiv (G_N M \omega)^{2/3}$, with $M \equiv m_1 + m_2$, to express r in terms of x . Once obtained the formula

$$\frac{G_N M}{r} = x \left[1 + \left(1 - \frac{\eta}{3} \right) x \right],$$

where $\eta \equiv \frac{m_1 m_2}{M^2}$, find the analog for $v \equiv \omega r$, which is

$$v^2 = x \left[1 + 2 \left(\frac{\eta}{3} - 1 \right) \right].$$

Exercise 22 *** Lorentz invariance in the non-relativistic limit**

Derive the non-relativistic limit of the Lorentz transformation

$$\begin{pmatrix} t' \\ \mathbf{x}' \end{pmatrix} = (1 - w^2)^{-1/2} \begin{pmatrix} 1 & -\mathbf{w} \\ -\mathbf{w} & 1 \end{pmatrix} \begin{pmatrix} t \\ \mathbf{x} \end{pmatrix}$$

Result:

$$\begin{aligned} \mathbf{x}_a &\rightarrow \mathbf{x}'_a = \mathbf{x} - \mathbf{w}t + \mathbf{v}_a(\mathbf{w} \cdot \mathbf{x}_a) \\ t &\rightarrow t' = t \end{aligned}$$

where \mathbf{w} is the boost velocity and $\mathbf{w}_a = \dot{\mathbf{x}}_a$. From the Lagrangian $L(x_a, v_a)$, let us define

$$\frac{\delta L}{\delta x_a^i} \equiv -\frac{d}{dt} \left(\frac{\partial L}{\partial v_a^i} \right) + \frac{\partial L}{\partial x_a}.$$

and on the equations of motion $\delta L / \delta x_a^i = 0$. Show that for *any* transformation $\delta x_a^i = x_a^{i'} - x_a^i$ one can write the Lagrangian variation as

$$\delta L = \frac{dQ}{dt} + \sum_a \frac{\delta L}{\delta x_a^i} \delta x_a^i + O(\delta x_a^2),$$

with

$$Q \equiv \sum_a \frac{\partial L}{\partial v_a^i} \delta x_a^i = \sum_a p_a^i \delta x_a^i.$$

If the transformation δx_a^i is a symmetry, the Lagrangian transforms as a total derivative, that is, in the case of boosts, $\delta L = w^i dZ^i / dt + O(w^2)$

for some function Z^i . Apply the above equation for δL in terms of Q to derive that invariance under boosts implies conservation of the quantity

$$G^i - \sum_a p_a^i t \equiv -Z^i + \sum_a x_a^i (\mathbf{p}_a \cdot \mathbf{v}_a) - \sum_a p_a^i t$$

and find the specific form of Z^i at Newtonian level and at 1PN level. Interpret G^i as the center of mass position, by imposing $G^i = 0$ and $\mathbf{x}_1 - \mathbf{x}_2 = \mathbf{r}$ find the expressions of $x_{1,2}^i$ in terms of the relative coordinate r^i .

Find the energy E of the system

$$E = \sum_a v_a^i p_a^i - L.$$

at Newtonian and 1PN level. Use the result of the previous step to express v_a in terms of the relative velocity v . Use the result of the previous exercise to find the energy of the circular orbit in terms of x and compare with the Schwarzschild result of ex. 8

Exercise 23 ***** Graviton loop

Diagrams like the one in fig. 8 do not affect the classical potential. How many powers of \hbar does this diagram contains? Estimate quantitatively its contribution to the effective action. Is it correct to expand the G_F in its quasi-static limit ($\omega^2 \ll k^2$)?

Exercise 24 ***** Field solution

Derive the expression of the potential of an isolated source at 1PN. Hint: you need to perform the path integral by setting an extra field in the integrand:

$$\langle h_{\mu\nu} \rangle = \int \mathcal{D}h_{\mu\nu} h_{\mu\nu} e^{iS_{eff}(J, h_{\mu\nu})} \quad (145)$$

which by virtue of eq. (115), is equivalent to taking a derivative with respect to the J source. Substitute the source-field coupling $J\Phi$ in eq. (118) with $J_\phi\phi + J_A A + J_\sigma\sigma$ given by $-m_a \int dt d^3x \delta(\mathbf{x} - X_a) [(1 + \frac{3}{2}v_a^2)\phi + A^i v_{ai} + \sigma^{ij} v_{aj} v_{ai}]$. Find the solution to the equation of motion of ϕ, A and σ_{ij} at Newtonian order by using

$$(\phi, A, \sigma_{ij}) = \frac{\delta Z_0[J]}{\delta (J_\phi, J_A, J_\sigma)}.$$

Use the presence of the term $c_d/2\partial_i\phi\partial_j\phi(\delta^{ik}\delta^{jl} + \delta^{il}\delta^{jk} - \delta^{ij}\delta^{kl})\sigma_{kl}$ in the Einstein-Hilbert action to derive $G_N m/r$ correction to the above solution.

Exercise 25 ***** 2PN potential of an isolated source

Similarly to the previous exercise, derive the 2PN potential of a static, isolated source. Note that in this case one can neglect the A and σ coupling in the world-line action, but *not* in the Einstein-Hilbert action.

Hint: According to the ansatz in eq. (128) ϕ couples to $T_{00} + T_{kk}$, σ_{ij} to T_{ij} and A_i to T_{0i} . Use the “bulk” 3-linear interaction, which can be deduced from the Einstein-Hilbert + gauge fixing:

$$S_{EH-h^3} \supset \int dt d^d x \frac{\phi}{\Lambda} \left[-4\dot{\phi}^2 + 4\dot{A}_i^2 + A_{i,j}A_{i,j} + \check{\sigma}_{ij}\check{\sigma}_{kl} \left(\delta_{ij}\delta_{kl} - \delta_{ik}\delta_{jl} - \delta_{il}\delta_{jk} \right) \right] - 8\phi\phi_k A_k$$

Result: one should derive the Schwarzschild metric in harmonic gauge

$$d\tau^2 = \frac{r - G_N M}{r + G_N M} dt^2 + \frac{(r + G_N M)^2}{r^2} \left[\delta_{ij} - \frac{x_i x_j}{r^2} \right] dx^i dx^j + \frac{r + G_N M}{r - G_N M} \frac{x_i x_j}{r^2} dx^i dx^j.$$

Exercise 26 ***** 2PN correction to the Energy momentum tensor of an isolated point particle

We now want to derive an expression for the “dressing” of the energy-momentum tensor because of the gravitational interaction. In quantum mechanics this correspond to evaluate the 1-point function of $T_{\mu\nu}$, that is $\langle T_{\mu\nu} \rangle$. In order to compute this it is useful to use the “background field method”, consisting in substituting in the integrand of the Feynman integral

$$h_{\mu\nu} \rightarrow h_{\mu\nu} + H_{\mu\nu},$$

perform the Gaussian integration and then keep only the terms linear in $H_{\mu\nu}$, which by definition describe the energy momentum tensor. Explain why this computation is the same as the one of the previous exercise, it just needs the use of 1 Green function less.

GW Radiation

In the previous chapters we have shown how to obtain an effective action describing the dynamics of a binary system at the orbital scale r in which gravitational degrees of freedom have been integrated out, resulting in a series expansion in v^2 , as in a conservative system odd powers of v are forbidden by invariance under time reversal.

The gravitational tensor in 3+1 dimensions has 6 physical degrees of freedom (10 independent entries of the symmetric rank 2 tensor in 3+1 dimensions minus 4 gauge choices): 4 of them are actually constrained, non radiative physical degrees of freedom, responsible for the gravitational potential, and the remaining 2 are radiative, or GWs.

In order to compute interesting observables, like the average energy flux emitted by or the radiation reaction on the binary system, we have to consider processes involving the emission of GW, i.e. on-shell gravitational modes escaping to infinity.

In order then to obtain an effective action for the sources alone, including the back-reaction for the emission of GWs, it will be useful to “integrate out” also the radiative degrees of freedom, with characteristic length scale $\lambda = r/v$, as it will be shown in the next sections. We aim now at writing the coupling of an extended source in terms of the energy momentum tensor $T^{\mu\nu}(t, x)$ moments, i.e. we want to introduce the *multipole expansion*. Here we use $T^{\mu\nu}$, as in ²⁵, to denote the term relating the effective action \mathcal{S}_{ext} relative to the external graviton emission

$$\mathcal{S}_{ext} = -\frac{1}{2} \int dt d^d x T^{\mu\nu}(t, x) h_{\mu\nu}(t, x), \quad (146)$$

to the gravitational mode generically denoted by $h_{\mu\nu}$. With this definition $T^{\mu\nu}$ receives contribution from both matter and the gravity *pseudo-tensor* appearing in the traditional GR description of the emission processes.

Given that the variation scale of the energy momentum tensor and of the radiation field are respectively r_{source} and λ , by Taylor-

²⁵ W. D. Goldberger and A. Ross. Gravitational radiative corrections from effective field theory. *Phys. Rev. D*, 81: 124015, 2010

expanding the standard term $T_{\mu\nu}h^{\mu\nu}$

$$\sum_n \frac{1}{n!} \partial_1 \dots \partial_n h_{\mu\nu}(t, x) \Big|_{x=0} \int d^d x' T^{\mu\nu}(t, x') x'^1 \dots x'^n, \quad (147)$$

we obtain a series in r_{source}/λ , which for binary systems gives

$$r_{source} = r \ll \lambda = r/v.$$

The results of the integral in eq. (147) are source moments that, following standard procedures not exclusive of the effective field theory approach described here, are traded for mass and velocity multipoles. For instance, the integrated moment of the energy momentum tensor can be traded for the mass quadrupole

$$Q^{ij}(t) \equiv \int d^d x T^{00}(t, x) x^i x^j, \quad (148)$$

by repeatedly using the equations of motion under the form $T^{\mu\nu}_{,\nu} = 0$:

$$\begin{aligned} \int d^d x \left[T^{0i} x^j + T^{0j} x^i \right] &= \int d^d x T^{0k} (x^i x^j)_{,k} \\ &= - \int d^d x T^{0k}_{,k} x^i x^j \\ &= \int d^d x \dot{T}^{00} x^i x^j = \dot{Q}^{ij} \end{aligned} \quad (149)$$

$$\begin{aligned} 2 \int d^d x T^{ij} &= \int d^d x \left[T^{ik} x^j_{,k} + T^{kj} x^i_{,k} \right] \\ &= \int d^d x \left[\dot{T}^{0i} x^j + \dot{T}^{0j} x^i \right] \\ &= \int d^d x \ddot{T}^{00} x^i x^j = \ddot{Q}^{ij}. \end{aligned} \quad (150)$$

The above equations also show that as for a composite binary system $T_{00} \sim O(v^0)$, then $T_{0i} \sim O(v^1)$ and $T_{ij} \sim O(v^2)$.

Taking as the source of GWs the composite binary system, the multipole series is an expansion in terms of $r/\lambda = v$, so when expressing the multipoles in terms of the parameter of the individual binary constituents, powers of v have to be tracked in order to arrange a consistent expansion. At lowest order in the multipole expansion and at v^0 order

$$\mathcal{S}_{ext}|_{v^0} = \frac{1}{\Lambda} \int dt d^d x T^{00}|_{v^0} \phi = \frac{M}{\Lambda} \int dt \phi, \quad (151)$$

where in the last passage the explicit expression

$$T^{00}(t, x)|_{v^0} = \sum_A m_A \delta^{(3)}(x - x_A(t)), \quad (152)$$

has been inserted. At order v the contribution from the first order derivative in ϕ have to be added the contribution of $T^{\mu\nu}|_v$, which gives

$$\mathcal{S}_{ext}|_v = -\frac{1}{\Lambda} \int dt d^d x \left(T^{00}|_{v^0} x^i \phi_{,i} + T^{0i}|_v A_i \right), \quad (153)$$

with

$$T^{0i}(t, x)|_v = \sum m_A v_{Ai} \delta^{(3)}(x - x_A(t)), \quad (154)$$

and neither T_{00} nor T_{ij} contain terms linear in v . Since the total mass appearing in eq. (151) is conserved (at this order) and given that in the center of mass frame $\sum_A m_A x_{Ai} = 0 = \sum_A m_A v_{Ai}$, there is no radiation up to order v . From order v^2 on, following a standard procedure, see e.g. ²⁶, it is useful to decompose the source coupling to the gravitational fields in irreducible representations of the $SO(3)$ rotation group, to obtain

$$\begin{aligned} S_{ext}|_{v^2}^{\mathbf{1}} &= -\frac{1}{2\Lambda} \int dt d^d x T^{0i}|_v x^j (A_{ij} - A_{ji}), \\ S_{ext}|_{v^2}^{\mathbf{0+2}} &= \frac{1}{4\Lambda} \int dt Q^{ij}|_{v^0} \left(\ddot{\phi}_{ij} - 2\phi_{,ij} - \frac{2}{d-2} \ddot{\phi} \delta_{ij} - \dot{A}_{i,j} - \dot{A}_{j,i} \right), \end{aligned} \quad (155)$$

were eqs. (149,150) and integration by parts have been used, $\mathbf{0}, \mathbf{1}, \mathbf{2}$ stand for the scalar, vector and symmetric-traceless representations of $SO(3)$, and

$$Q^{ij}|_{v^n} = \int d^d x T^{00}|_{v^n} x^i x^j. \quad (156)$$

The $\mathbf{1}$ part matches the second term in eq. (157), and it is not responsible for radiation as it couples A_i to the conserved angular momentum. Let us pause for a moment to overlook our procedure and identify the $\mathbf{0}$ and $\mathbf{2}$ terms. We want to describe a composite system, the coalescing binary made of two compact objects. At the scale of GW radiation $\lambda = r/v \gg r$ the binary system can be seen as a single object characterized by its coupling to the background gravitational fields. The point action (104) describe point particles, but extended objects experiences tidal field due to their non-zero size. It is possible to parametrize such effects by effective terms that take into account finite size in a completely general way, via the expression

$$S_{ext} \supset \int d\tau \left(-M - \frac{1}{2} S_{ab} \omega_\mu^{ab} u^\mu + \frac{c_Q}{2} I^{ij} E_{ij} + \frac{c_J}{2} J^{ij} B_{ij} + \frac{c_O}{2} I^{ijk} \partial_i E_{jk} + \dots \right) \quad (157)$$

where ω_μ^{ab} is the spin connection coupling to the total angular momentum, while the electric (magnetic) tensor E_{ij} (B_{ij}) is defined by

$$\begin{aligned} E_{ij} &= C_{\mu\nu j} u^\mu u^\nu, \\ B_{ij} &= -\frac{1}{2} \epsilon_{i\mu\nu\rho} u^\rho C^{\mu\nu}_{j\sigma} u^\sigma, \end{aligned} \quad (158)$$

decomposing the Weyl tensor $C_{\mu\alpha\nu\beta}$ analogously to the electric and magnetic decomposition of the standard electromagnetic tensor $F_{\mu\nu}$. The $\mathbf{0} + \mathbf{2}$ term in eq. (155) reproduces at linear order the coupling

²⁶ M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008

$Q^{ij}R^0_{ij}$ term in eq. (157), allowing to identify I_{ij} with Q_{ij} at linear order and to impose $c_Q = 1$.

This amounts to decompose the source motion in terms of the world-line of its center of mass and moments describing its internal dynamics. The I^{ij} , I^{ijk} , J^{ij} tensors are the lowest order in an infinite series of source moments, the 2^{nth} electric (magnetic) moment in the above action scale at leading order as mr_{source}^n (mvr_{source}^n), and they couple to the Taylor expanded E_{ij} (B_{ij}) which scales as $L^{-(1+n)}$, showing that the above multipole expansion is an expansion in terms of r_{source}/L . Note that the world-line of the composite object cannot couple to R and $R_{\mu\nu}$, as by the Einstein equation they vanish where sources are not present, thus such couplings can be neglected if we are interested in describing how a composite object respond to the gravity field of other objects.²⁷ As linear terms in the Ricci tensor and Ricci scalar cannot appear, the terms involving the least number of derivatives are the ones written above in eq. (157), in terms of the (traceless part of the) Riemann tensor. Out of the Riemann tensor, we select only its trace-free part (10 components out of 20 in $3+1$ dimensions, which are then re-arranged in the 5 components of the electric Weyl tensor E_{ij} and 5 components in the magnetic Weyl tensor B_{ij}).

Note that the multipoles, beside being intrinsic, can also be induced by the tidal gravitational field or by the intrinsic angular momentum (spin) of the source. For quadrupole moments the tidal induced quadrupole moments $I_{ij}, J_{ij}|_{tidal} \propto E_{ij}, B_{ij}$ give rise to the following terms in the effective action

$$\mathcal{S}_{tidal} = \int d\tau \left[c_E E_{ij} E^{ij} + c_B B_{ij} B^{ij} \right]. \quad (159)$$

This is also in full analogy with electromagnetism, where for instance particles with no permanent electric dipole experience a quadratic coupling to an external electric field. Eq. (159) can be used to describe a single, spin-less compact object in the field of its binary system companion. Considering that the Riemann tensor generated at a distance r by a source of mass m goes as m/r^3 , the finite size effect given by the $E_{ij}E^{ij}$ term goes as $c_E m^2/r^6$. For dimensional reasons $c_E \sim G_N r_{source}^5$ ²⁸, thus showing that the finite size effects of a spherical symmetric body in the binary potential are $O(Gm/r)^5$ times the Newtonian potential, a well known result which goes under the name of *effacement principle*²⁹ (the coefficient c_E actually vanishes for black holes in $3+1$ dimensions³⁰).

In order to simplify the calculation, we work from now on in the transverse-traceless (TT) gauge, in which the only relevant radiation field is the traceless and transverse part of σ_{ij} . The presence of the other gravity polarizations is required by gauge invariance.

²⁷ Equivalently, it can be shown that the field redefinition $g_{\mu\nu} \rightarrow g_{\mu\nu} + \delta g_{\mu\nu}$ with

$$\delta g_{\mu\nu} = \int d\tau \frac{\delta(x^\alpha - x^\alpha(\tau))}{\sqrt{-g}} \left[\left(-c_R + \frac{c_V}{2} \right) g_{\mu\nu} - c_V u_\mu u_\nu \right]$$

can be used to set to zero the above terms linear in the curvature, see for details.

W. D. Goldberger. Les houches lectures on effective field theories and gravitational radiation. In *Les Houches Summer School - Session 86: Particle Physics and Cosmology: The Fabric of Spacetime*, 2007

²⁸ Walter D. Goldberger and Ira Z. Rothstein. An Effective field theory of gravity for extended objects. *Phys.Rev.*, D73:104029, 2006

²⁹ T. Damour. *Gravitational radiation and the motion of compact bodies*, pages 59–144. North-Holland, Amsterdam, 1983b

³⁰ T. Damour. Gravitational radiation and the motion of compact bodies. In N. Deruelle and T. Piran, editors, *Gravitational Radiation*, pages 59–144. North-Holland, Amsterdam, 1983a; and Barak Kol and Michael Smolkin. Black hole stereotyping: Induced gravitostatic polarization. *JHEP*, 1202:010, 2012. DOI: 10.1007/JHEP02(2012)010

Discarding all fields but the TT-part of the σ_{ij} field, at order v^3 one has

$$S_{ext}|_{v^3} = \frac{1}{2} \int dt d^d x T^{ij}|_{v^2} x^k \sigma_{ij,k} \quad (160)$$

and using the decomposition see ex. 28

$$\int d^d x T^{ij} x^k = \frac{1}{6} \int d^d x \ddot{T}^{00} x^i x^j x^k + \frac{1}{3} \int d^d x \left(\dot{T}^{0i} x^j x^k + \dot{T}^{0j} x^i x^k - 2\dot{T}^{0k} x^i x^j \right), \quad (161)$$

we can re-write, see ex.29,

$$S_{ext}|_{v^3} = \int dt \left(\frac{1}{6} Q^{ijk} E_{ij,k} - \frac{2}{3} P^{ij} B_{ij} \right) \quad (162)$$

where

$$Q^{ijk} = \int d^d x T^{00} x^i x^j x^k, \quad (163)$$

and

$$P^{ij} = \int d^d x \left(\epsilon^{ikl} T_{kl}^0 x_l x^j + \epsilon^{jkl} T_{kl}^0 x_l x^i \right), \quad (164)$$

allowing to identify $J^{ij} \leftrightarrow P^{ij}$ and $I^{ijk} \leftrightarrow Q^{ijk}$ at leading order, with $c_J = -4/3$ and $c_O = 1/3$. In order to derive eq. (164) see ex. 29. Note that from now on it is understood that the GW field is the TT one, so it couple only to the trace-free part of the source

At v^4 order the $T^{ij} x^k x^l \sigma_{ij,kl}$ term, beside giving the leading hexadecapole term (or 2^{4th} -pole) and v corrections to the leading magnetic quadrupole and electric octupole, also gives a v^2 correction to the leading quadrupole interaction $I^{ij} E_{ij}$, which can be written as (we give no demonstration here, see sec. 3.5 of ³¹ for the general method)

$$S_{ext}|_{v^4} \supset \int dt d^3 x \left[T^{00}|_{v^2} + T^{kk}|_{v^2} - \frac{4}{3} \dot{T}^{0k}|_v x^k + \frac{11}{42} \ddot{T}_{00}|_{v^0} x^2 \right] \times \left(x^i x^j - \frac{\delta^{ij}}{d} x^2 \right) E_{ij}. \quad (165)$$

For the systematics at higher orders we refer to the standard textbook ³².

Matching between the radiation and the orbital scale

In the previous subsection we have spelled out the general expression of the effective multipole moments in terms of the energy-momentum tensor moments. However we have only used two ingredients from the specific binary problem

- $T_{00} \sim mv^0$

³¹ M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008


³² M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008

- the source size is r and the length variation of the background is $\lambda \sim r/v$.

Now we are going to match the coefficients appearing in eq. (157) with the parameters of the specific theory at the orbital scale.

At leading order $Q_{ij}|_{v^0} = \sum_A m_A x_{Ai} x_{Aj} = \mu r^i r^j$ and the v^2 corrections to T_{00} can be read from diagrams in figs. 10. As ϕ couples to $T_{00} + T_{kk}$ and σ_{ij} to T_{ij} , from the diagrams one obtains³³

$$\begin{aligned} \int d^3x (T_{00} + T_{kk}) \Big|_{v^2} &= \sum_A \frac{3}{2} m_A v_A^2 - \sum_{B \neq A} \frac{G_N m_A m_B}{r}, \\ \int d^3x T_{kk} \Big|_{v^2} &= \sum_A \frac{1}{2} m_A v_A^2 - \frac{1}{2} \sum_{B \neq A} \frac{G_N m_A m_B}{r}. \end{aligned} \quad (166)$$



³³ Note that since only $\int T_{kk}$ is needed, and not T_{kk} itself, it could have been computed from eq. (150) instead of from the diagram in fig. 10b.

Figure 10: Graph dressing T_{00} at v^2 order and T_{ij} at leading order. The external radiation graviton does not carry momentum but it is Taylor expanded.

These diagrams are obtained by performing the Gaussian path integral with an external gravitational mode $H_{\mu\nu}$ on which one does not have to integrate over. This correspond to adopting the “background-field method”, in which every graviton in the original Lagrangian is splitted according to $h_{\mu\nu} \rightarrow h_{\mu\nu} + H_{\mu\nu}$ and then integration is performed on $h_{\mu\nu}$ alone. The term linear in $H_{\mu\nu}$ in the resulting effective action is by definition (proportional to) the energy-momentum tensor.

Integrating out the radiating graviton and mass renormalization




Figure 11: Diagram giving the leading term of the amplitude describing radiation back-reaction on the sources.

We have now built an effective theory for extended objects in terms of the source moments and we also shown how to match the




Figure 12: Next-to-leading order term in the back-reaction amplitude.

orbital scale with the theory describing two point particles experiencing mutual gravitational attraction. We can further use the extended object action in eq. (157) to integrate out the gravitational radiation to obtain an effective action S_{mult} for the source multipoles alone.

In order to perform such computation, boundary conditions *asymmetric in time* have to be imposed, as no incoming radiation at past infinity is required. Using the standard *Feynman* propagator, which ensures a pure in-(out-)going wave at past (future) infinity, would lead to a non-causal evolution as it can be shown by looking at the following toy model³⁴, which is defined by a scalar field Ψ coupled to a source J :

$$S_{toy} = \int d^{d+1}x \left[-\frac{1}{2} (\partial\psi)^2 + \psi J \right]. \quad (167)$$

We may recover the field generated by the source J as

$$\psi(t, x) = \int d^{d+1}x' G_F(t - t', x - x') J(t', x'). \quad (168)$$

In a causal theory ψ would be given by the same eq. (168) but with the Feynman propagator replaced by the retarded one $G_{Ret}(t, x)$, however it is not possible to naively use the retarded propagator in the action (167), as it would still yield non-causal equations of motions as

$$\begin{aligned} & \int dt d^3x dt' d^3x' \psi(t, x) G_{ret}^{-1}(t - t', x - x') \psi(t', x') \\ &= \frac{1}{2} \int dt d^3x dt' d^3x' \psi(t, x) \left(G_{ret}^{-1}(t - t', x - x') + G_{adv}^{-1}(t - t', x - x') \right) \psi(t', x') \end{aligned} \quad (169)$$

This problem was not present in the conservative dynamics described in sec. as there we had a closed system with no leak of energy: there we actually had the Feynman Green function which is symmetric in its arguments, see ex. 5.

However there is a consistent way to define an action for non-conservative system with asymmetric time boundary condition: by adopting a generalization of the Hamilton's variational principle similar to the closed-time-path, or in-in formalism (first proposed in

³⁴ M. Tiglio C. R. Galley. Radiation reaction and gravitational waves in the effective field theory approach. *Phys. Rev., D79:124027*, 2009

³⁵, see ³⁶ for a review) as described in ³⁷, which requires a *doubling* of the field variables. For instance the toy model in eq. (167) is modified so that the generating functional for connected correlation functions in the in-in formalism has the path integral representation

$$e^{iS_{eff}[J_1, J_2]} = \int \mathcal{D}\psi_1 \mathcal{D}\psi_2 \times \exp \left\{ i \int d^{3+1}x \left[-\frac{1}{2}(\partial\psi_1)^2 + \frac{1}{2}(\partial\psi_2)^2 - J_1\psi_2 + J_2\psi_1 \right] \right\}. \quad (170)$$

In this toy example the path integral can be performed exactly, and using the Keldysh representation ³⁸ defined by $\Psi_- \equiv \Psi_1 - \Psi_2$, $\Psi_+ \equiv (\Psi_1 + \Psi_2)/2$, one can write

$$S_{eff}[J_+, J_-] = \frac{i}{2} \int d^{d+1}x d^{d+1}y J_B(x) G^{BC}(x-y) J_C(y), \quad (171)$$

where the B, C indices take values $\{+, -\}$ and

$$G^{BC}(t, \mathbf{x}) = \begin{pmatrix} 0 & G_{adv}(t, \mathbf{x}) \\ G_{ret}(t, \mathbf{x}) & 0 \end{pmatrix}. \quad (172)$$

In our case, the lowest order expression of the quadrupole in terms of the binary constituents world-lines x_A , i.e.

$$Q_{ij}|_{v^0} = \sum_{A=1}^2 m_A \left(x_{Ai} x_{Aj} - \frac{\delta_{ij}}{d} x_{Ak} x_{Ak} \right), \quad (173)$$

is doubled to

$$\begin{aligned} Q_{-ij}|_{v^0} &= \sum_{A=1}^2 m_A \left[x_{-Ai} x_{+Aj} + x_{+Ai} x_{-Aj} - \frac{2}{d} \delta_{ij} x_{+Ak} x_{-Ak} \right] \\ Q_{+ij}|_{v^0} &= \sum_{A=1}^2 m_A \left[x_{+Ai} x_{+Aj} - \frac{1}{d} \delta_{ij} x_{+A}^2 + O(x_-^2) \right]. \end{aligned} \quad (174)$$

The word-line equations of motion that properly include radiation reaction effects are given by

$$0 = \frac{\delta S_{eff}[x_{1\pm}, x_{2\pm}]}{\delta x_{A-}} \Bigg|_{\substack{x_{A-}=0 \\ x_{A+}=x_A}}. \quad (175)$$

At lowest order, by integrating out the radiation graviton, i.e. by computing the diagram in fig. 11, one obtains the Burke-Thorne ³⁹ potential term in the effective action S_{mult}

$$\begin{aligned} iS_{mult}|_{fig. 11} &= i^2 \int dt dt' Q_{-ij}(t) Q_{+kl}(t') E_{+ij}(t) E_{-kl}(t') \\ &= \frac{-i^3}{\Lambda^2} \int dt dt' Q_{-ij}(t) Q_{+kl}(t') \int_{\mathbf{k}} \frac{d\omega}{2\pi} \frac{e^{i\omega(t-t')}}{k^2 - (\omega + i\epsilon)^2} \\ &\times \left[\frac{\omega^4}{8} (\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk} - 2\delta_{ij} \delta_{kl}) \right. \\ &+ \frac{\omega^2}{8} (k_i k_k \delta_{jl} + k_i k_l \delta_{jk} + k_j k_k \delta_{il} + k_j k_l \delta_{ik}) \\ &\left. - \frac{1}{8} (\omega^4 \delta_{ij} \delta_{kl} + \omega^2 \delta_{ij} k_k k_l + \omega^2 k_i k_j \delta_{kl} + k_i k_j k_k k_l) \right] \end{aligned} \quad (176)$$

³⁵ J. S. Schwinger. Brownian motion of a quantum oscillator. *J. Math. Phys.*, 2: 407–432, 1961

³⁶ B. DeWitt. Effective action for expectation values. In R. Penrose and C. J. Isham, editors, *Quantum concepts in Space and Time*. Clarendon Press, Oxford, 1986

³⁷ C. R. Galley. The classical mechanics of non-conservative systems. *Phys. Rev. Lett.*, 110:174301, 2013

³⁸ L. V. Keldysh. Diagram technique for nonequilibrium processes. *Zh. Eksp. Teor. Fiz.*, 47:1515–1527, 1964

³⁹ W. L. Burke and K. S. Thorne. In S. I. Fickler M. Carmeli and L. Witten, editors, *Relativity*, pages 209–228. Plenum, New York, 1970

After performing the angular integration

$$\begin{aligned}\int d\Omega k_i k_j &= \frac{4}{3}\pi k^2 \delta_{ij}, \\ \int d\Omega k_i k_j k_k k_l &= \frac{4}{15}\pi k^4 \left(\delta_{ij}\delta_{kl} + \delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk} \right)\end{aligned}\quad (177)$$

and using

$$\int_0^\infty dk \frac{k^a}{k^2 - (\omega + i\epsilon)^2} = \frac{\pi}{\cos(a\frac{\pi}{2})} \left[\frac{1}{-(\omega + i\epsilon)^2} \right]^{\frac{1-a}{2}}, \quad (178)$$

one finds the result

$$iS_{mult}|_{fig. 11} = i\frac{G_N}{5} \int_{-\infty}^\infty \frac{d\omega}{2\pi} \omega^5 Q_{-ij}(\omega) Q_{+ij}(\omega) \quad (179)$$

$$= \frac{G_N}{5} \int dt Q_{-ij}(t) \frac{d^6 Q_{+ij}(t)}{dt^6} \quad (180)$$

and using eq. (175) the equation of motion follows:

$$\ddot{x}_i|_{fig. 11} = -\frac{2}{5} Q_{ij}^{(5)} x^j \quad (181)$$

Corrections to the leading effect appears when considering as in the previous subsection higher orders in the multipole expansion: the 1PN correction to the Burke Thorne potential were originally computed in ⁴⁰ and re-derived with effective field theory methods in ⁴¹.

The genuinely non-linear effect, computed originally in ⁴² and within effective field theory methods in ⁴³, appears at relative 1.5PN order and it is due to the diagram in fig. 12. The contribution to the effective action can be obtained by considering the contribution respectively from the $\sigma^2\psi$, $\sigma\phi A$, $\sigma\phi^2$, $A^2\phi$, $A\phi^2$, ϕ^3 vertices, which give the effective action contribution

⁴⁰ B. R. Iyer and C. M. Will. Postnewtonian gravitational radiation reaction for two-body systems. *Phys. Rev. Lett.*, 70: 113, 1993

⁴¹ C. R. Galley and A. K. Leibovich. Radiation reaction at 3.5 post-newtonian order in effective field theory. *Phys. Rev. D*, 86:044029, 2012

⁴² Luc Blanchet and Thibault Damour. Tail transported temporal correlations in the dynamics of a gravitating system. *Phys.Rev.*, D37:1410, 1988; and L. Blanchet. Time asymmetric structure of gravitational radiation. *Phys.Rev.*, D47: 4392-4420, 1993

⁴³ S. Foffa and R. Sturani. Tail terms in gravitational radiation reaction via effective field theory. *Phys. Rev. D*, 87: 044056, 2013

$$\begin{aligned}
iS_{eff}|_{fig. 11} = & -i \frac{M}{16\Lambda^4} \int \frac{d\omega}{2\pi} (Q_{-ij}(\omega) Q_{+kl}(-\omega) + I_{+ij}(-\omega) I_{-kl}(\omega)) \\
& \int \frac{d^d k}{(2\pi)^d} \frac{d^d q}{(2\pi)^d} \frac{1}{\mathbf{k}^2 - (\omega - ia)^2} \frac{1}{(\mathbf{k} + \mathbf{q})^2 - (\omega - ia)^2} \frac{1}{\mathbf{q}^2} \\
& \left\{ -\frac{1}{8} \omega^6 \left(\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk} - \frac{2}{d-2} \delta_{ij} \delta_{kl} \right) \right. \\
& + \frac{1}{2} \omega^4 (q_i q_k \delta_{jl}) \\
& + \frac{1}{2c_d} \omega^2 \left(-\frac{k_\delta^2 q_i q_j}{d-2} \delta_{kl} - q_i q_j k_k k_l - 2k_i q_j k_k q_l - 3q_i q_j q_k k_l - q_i q_j q_k q_l \right) \\
& + \frac{1}{2} \omega \left[k_i k_j q_k q_l - q_i k_j k_k q_l + \delta_{ik} (k_j k_l \mathbf{k}^2 + k_j q_l (\mathbf{k} \mathbf{q}) + \mathbf{k}^2 k_j q_l + \mathbf{k} \mathbf{q} k_j k_l) \right] + \\
& \frac{1}{2c_d} \omega \left(2q_i k_j q_k k_l - k_i k_j q_k q_l + q_i k_j q_k q_l - k_0^2 \delta_{kl} \frac{q_i q_j}{d-2} \right) \\
& - \frac{1}{2c_d} \omega \left[k_i k_j k_k k_l + k_i k_j q_k q_l + 2k_i k_j k_k q_l + \right. \\
& \left. \frac{\omega^2}{d-2} \left(\delta_{ij} k_k k_l + \delta_{kl} k_i k_j + \delta_{ij} q_k q_l + 2\delta_{ij} k_k q_l + \frac{\omega}{d-2} \delta_{ij} \delta_{kl} \right) \right] \left. \right\}. \quad (182)
\end{aligned}$$

where we have been careful to work in generic d spatial dimensions: this diagram has a logarithmic divergence in $d = 3$, which is cured by performing the computation in generic d dimension. In order to isolate the cause of the divergence, let us focus on the first line of eq. (182) within the curly bracket, which is the simplest integral to perform, has it boils down to integrate the two propagators of the previous line. After using (see eq.(8-7) of ⁴⁴)

$$\begin{aligned}
\int_{\mathbf{k}} \frac{1}{[k^2 - \omega_1^2]^{2a}} \frac{1}{[(\mathbf{k} + \mathbf{q})^2 - \omega_2^2]^{2b}} &= \frac{1}{(4\pi)^d/2} \frac{\Gamma(a+b-d/2)}{\Gamma(a)\Gamma(b)} \\
\times \int_0^1 dx x^{a-1} (1-x)^{b-1} &\left[x(1-x)q^2 - x\omega_1^2 - (1-x)\omega_2^2 \right]^{d/2-a-b} \quad (183)
\end{aligned}$$

and the identity

$$\int_0^1 x^a (1-x)^b = \frac{\Gamma(1+a)\Gamma(1+b)}{\Gamma(2+a+b)}, \quad (184)$$

we get a term proportional to

$$\int_{\mathbf{q}} \frac{1}{q^2} \left[q^2 - \frac{(\omega + i\epsilon)^2}{x(1-x)} \right]^{d/2-2} \quad (185)$$

which diverges logarithmically as $q \rightarrow \infty$ for $d = 3$. Performing the full computation, whose details can be found in ⁴⁵, one obtains

$$\begin{aligned}
S_{eff}|_{fig.12} = & -\frac{1}{5} G_N^2 M \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \omega^6 \left(\frac{1}{d-3} - \frac{41}{30} + i\pi \text{sgn}(\omega) - \log \pi + \gamma \right) \\
& \times \left(\frac{\omega^2}{\mu^2} \right) \times \left(Q_{ij-}(\omega) Q_{ij+}(-\omega) + Q_{ij-}(-\omega) Q_{ij+}(\omega) \right). \quad (186)
\end{aligned}$$

⁴⁴ C. Itzinkinson and J. B. Zuber. *Quantum field theory*. Mac Graw-Hill International Book Company, 1980

⁴⁵ S. Foffa and R. Sturani. Tail terms in gravitational radiation reaction via effective field theory. *Phys. Rev. D*, 87: 044038, 2013

Note the appearance of the arbitrary scale μ (with dimension of inverse-length) because in generic d dimensions we have $\Lambda^{-2} = 32\pi G_N \mu^{3-d}$, with μ setting the scale of the d -dimensional Newton's constant.

Let us look at the physical process we are considering in order to interpret this result: a GW is emitted from the quadrupole, scattered by the curvature and absorbed by the binary itself. In the limit of very large q , i.e. when the scattering occurs arbitrarily close to the binary system, this amplitude diverges. We cannot trust our effective theory arbitrarily close to the source, where the system cannot be described as a single object with a quadrupole, but we have to match with the fundamental theory of a binary system. The finite numbers we get with the result are then meaningless, as they are obtained by pushing our effective theory farther than it can actually be trusted. However the logarithmic terms *is* physical. By cutting the \mathbf{q} integral at any finite scale q_0 , we would have obtained a term $\log(\omega/q_0)$, which would tell that integrating out gravitational interactions up to an arbitrary finite scale, will introduce a logarithmic scaling of the result. Of course it does not make sense that a *physical* result depends on an arbitrary scale, indeed shifting μ to a new value μ' would change the *unphysical* finite terms in eq. (186) by a finite amount proportional to $\log(\mu'/\mu)$. How to make physical sense out of this result? The source of the Newtonian potential has a logarithmic dependence of the scale μ : it changes if we probe it at different distances, because of the "cloud" of GW which is continuously emitted and absorbed by the binary systems, through their quadrupole coupling to the source.

It is possible to derive the correct finite terms to complete the result by using the correct theory to describe the physics very close to the binary system, which at short distances cannot be treated as a point of zero size, but rather as a binary system with size $0 < r \ll \lambda r/v$.

We note the presence of the logarithmic term which is non-analytic in ω -space and non-local (but causal) in direct space: after integrating out a mass-less propagating degree of freedom the effective action is not expected to be local ⁴⁶.

In order to make sense of the result (186) we have first to regularize it, which can be done by adding a *local* counter term to the action (157) given by M_{ct} defined by

$$M_{ct} = -\frac{2G_N^2}{5} M \left(\frac{1}{\epsilon} + \gamma - \log \pi \right) Q_{-ij} Q_{+ij}^{(6)}. \quad (187)$$

The combination of the *bare* monopole mass term M in eq.(157) plus the above M_{ct} give the mass which is the measured parameter.

⁴⁶ T. Appelquist and J. Carazzone. Infrared singularities and massive fields. *Phys. Rev. D*, 11:2856, 1975

However, this is not the full story, as we still have to make sense of the logarithm appearing in eq.(186).

The scale μ appearing in the logarithm is arbitrary, but the presence of the logarithm is indeed *physical*: the way to avoid an unphysical dependence on μ of the physical effective action is to *assume* that M is μ dependent, that is the monopole mass of the binary system depends on the scale at which we are probing it, see ex. 31 for a thorough way to derive how the mass of the binary system is affected by the GW emission.

According to the standard renormalization procedure, one can define a renormalized mass $M^{(R)}(\mu)$ for the monopole term in the action (157), depending on the arbitrary scale μ in such a way that physical quantities (like the force derived from the effective action of the composite system) will be μ -independent. The force looks like

$$\delta\ddot{x}_{Ai}(t)|_{\log} = -\frac{8}{5}x_{aj}(t)G_N^2M \int_{-\infty}^t dt' Q_{ij}^{(7)}(t') \log[(t-t')\mu], \quad (188)$$

and it can be separated into a t -dependent and a t -independent part, with the t -independent part being:

$$\mu \frac{d\delta\ddot{x}_{Ai}(t)}{d\mu} = \frac{8}{5}x_{aj}(t)G_N^2M \int_{-\infty}^t dt' Q_{ij}^{(7)}(t') = \frac{8}{5}x_{aj}(t)G_N^2MQ^{(6)}(t). \quad (189)$$

We can see how this effect gives a logarithmic shift δM to the monopole mass:

$$\frac{d\delta M}{dt} = -\sum_A m_A \delta\ddot{x}_{Ai} \dot{x}_{Ai}. \quad (190)$$

Substituting eq. (188) into eq. (190) and using the leading order quadrupole moment expression in eq. (173) allows to turn the right hand side of eq. (190) into a total time derivative, enabling to identify the logarithmic mass shift as ⁴⁷

$$\mu \frac{d\delta M^{(R)}}{d\mu} = -\frac{2G_N^2M}{5} \left(2Q_{ij}^{(5)}Q_{ij}^{(1)} - 2Q_{ij}^{(4)}Q_{ij}^{(2)} + Q_{ij}^{(3)}Q_{ij}^{(3)} \right). \quad (191)$$

This classical renormalization of the mass monopole term (which can be identified with the Bondi mass of the binary system, that does not include the energy radiated to infinity) is due to the fact that the emitted *physical* (not virtual!) radiation is scattered by the curved space and then absorbed, hence observers at different distance from the source would not agree on the value of the mass.

The ultraviolet nature of the divergence points to the incompleteness of the effective theory in terms of multipole moments: the terms analytic in ω in eq. (186) are sensitive to the short distance physics and their actual value should be obtained by going to the theory at orbital radius.

⁴⁷ L. Blanchet, S. L. Detweiler, A. Le Tiec, and B. F. Whiting. High-order post-newtonian fit of the gravitational self-force for circular orbits in the schwarzschild geometry. *Phys. Rev.*, D81, 2010

Emitted flux

We have now shown how to perform the matching between the theory of extended objects with multipoles and the theory at the orbital scale. Taking the action for extended bodies in eq. (157) as a starting point, the emitted GW-form and the total radiated power can be computed in terms of the source multipoles by using Feynman diagrams with one external radiating gravitational particle. Following the same rules given at the end of the previous chapter for the computation of the effective energy momentum tensor and the effective potential, we can write here the effective action for a composite particle described by the action (157). At leading order the effective action describing the emission/absorption of a GW described in fig. 13, one does not need to perform any computation and it can be just read from the action (157)

$$iS_{eff}(M, Q, \dots, \sigma_{ij}) \supset i \int dt \ddot{Q}_{ij} \frac{\sigma_{ij}}{4\Lambda} \quad (192)$$

The GW-form can be computed by convolving the appropriate Green function with the source, which at this order is simply \ddot{Q}_{ij} and it is then possible to derive the actual GW-form

$$\begin{aligned} \sigma_{ij}(t, x) \supset & 8\pi G_N \Lambda_{ij;kl} \int dt' d^d x' G_R(t - t', x - x') \\ & \times \left[\ddot{I}_{kl} + \frac{4}{3} \epsilon_{lmn} \dot{J}_{mk,n} - \frac{1}{3} \ddot{I}_{klm,m} \right], \end{aligned} \quad (193)$$

where $\Lambda_{ij,kl}$ is the TT-projector $\Lambda_{ij;kl}$ defined in eq. (22).




Figure 13: Diagram representing the emission of a GW from a quadrupole source

Analogously to what shown in the previous subsection, we have to take into account the GW interaction with the space time curvature produced by the source itself. Including such effect give rise to a *tail* effect, accounted by the diagram in fig. 14, which gives a contribution to the GW amplitude and phase ⁴⁸

$$\begin{aligned} \sigma_{ij}|_{fig. 14} \supset & \Lambda_{ij;kl} \pi M G \int_{\mathbf{k}} \frac{d\omega}{2\pi} \int_{\mathbf{k}} e^{-i\omega t + i\mathbf{k}\mathbf{x}} \omega^4 \\ & \frac{1}{k^2 - (\omega + i\epsilon)^2} \int_{\mathbf{q}} \frac{q^{-2}}{(\mathbf{k} + \mathbf{q})^2 - (\omega + i\epsilon)^2} I_{kl}(\omega) \end{aligned} \quad (194)$$

⁴⁸ L. Blanchet and Gerhard Schaefer. Gravitational wave tails and binary star systems. *Class.Quant.Grav.*, 10:2699–2721, 1993; and A. Ross R. A. Porto and I. Z. Rothstein. Spin induced multipole moments for the gravitational wave amplitude from binary inspirals to 2.5 post-newtonian order. *JCAP*, 1209:028, 2012




Figure 14: Emission of a GW from a quadrupole source with post-Minkowskian correction represented by the scattering off the background curved by the presence of binary system.

The solution of the above integral is a bit involved, now we just want to underline that it contains an *infra-red* divergence for $q \rightarrow 0$: performing first the integral over ω yields a term proportional to

$$\int_{\mathbf{q}} \frac{1}{q^2} \frac{1}{2\mathbf{k}\mathbf{q}}. \quad (195)$$

The infra-red singularity in the phase of the emitted wave is un-physical as it can be absorbed in a re-definition of time in eq. (194). Moreover any experiment, like LIGO and Virgo for instance, can only probe phase *differences* (e.g. the GW phase difference between the instants when the wave enters and exits the experiment sensitive band) and the un-physical dependencies on the regulator ϵ and on the subtraction scale μ drops out of any observable.

The total emitted flux can be computed once the amplitude of the GW has been evaluated, via the standard formula, see eq. (71)

$$P = \frac{r^2}{32\pi G_N} \int d\Omega \langle \dot{h}_{ij} \dot{h}_{ij} \rangle, \quad (196)$$

but there is actually a shortcut, as the emission energy rate can be computed directly from the amplitude effective action (192) without solving for σ_{ij} . The derivation of the shortcut formula requires the interpretation of

$$A_h \equiv -k^2 Q_{ij} \frac{\epsilon_{ij}(\mathbf{k})}{4\Lambda} \quad (197)$$

as a *probability amplitude* for emitting a GW with polarization tensor $\epsilon_{ij}(\mathbf{k}, h)$ (the polarization tensor). The differential probability per unit of time for emitting a GW over the full phase space volume is indeed given by

$$dP(k) = \frac{1}{2T} \frac{d^3k}{(2\pi)^3} |k^2 Q_{ij}(k) \epsilon_{ij}(\mathbf{k})|^2. \quad (198)$$

This formula is usually obtained in quantum mechanics by applying the *optical theorem*, whose demonstration is obtained by noting that $|A_h|$ is obtained by taking the imaginary part of eq. (176), with the retarded propagator replaced by the Feynman one and using

$$\frac{1}{k^2 - \omega^2 + i\epsilon} = P \frac{1}{k^2 - \omega^2} + i\pi\delta(k^2 - \omega^2). \quad (199)$$

The imaginary part is then interpreted as a *probability loss* per unit of k space, as eq. 176 can be rewritten as

$$\begin{aligned} \text{Im } S'_{mult}|_{fig. 11} &= \int_{\mathbf{k}} \frac{d\omega}{2\pi} \sum_h |A_h(\omega, \mathbf{k})|^2 \pi\delta(k^2 - \omega^2) \\ &= \int_{\mathbf{k}} \frac{1}{2k} \sum_h |A_h(k, \mathbf{k})|^2, \end{aligned} \quad (200)$$

where \sum_h indicates sum over the two GW elicities. The imaginary part of the action gives the integrated probability loss, which weighted by one power of k and divided by the total observation time, gives the average energy flux. The last missing ingredient to get to the final formula is the sum over the two polarizations, which depends on \mathbf{k} via the specific form of the Λ projector. For the electric quadrupole term we have

$$\begin{aligned} &\int \frac{d\Omega}{4\pi} \sum_h \epsilon_{ij}^*(\mathbf{k}) \epsilon_{kl}(\mathbf{k}) \\ &= \int \frac{d\Omega}{4\pi} \frac{1}{2} \left[\delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk} - \delta_{ik}n_jn_l - \delta_{jl}n_in_k - \delta_{il}n_jn_k - \delta_{jk}n_in_k + n_in_jn_kn_l \right. \\ &\quad \left. - \delta_{ij}\delta_{kl} + \delta_{ij}n_kn_l + \delta_{kl}n_in_j \right] \\ &= \frac{1}{5} \left(\delta_{ik}\delta_{jl} + \delta_{ik}\delta_{jl} - \frac{8}{3}\delta_{ij}\delta_{kl} \right). \end{aligned} \quad (201)$$

For the magnetic quadrupole and electric octupole see ex. 36.

After summing over GW polarizations one gets:

$$\begin{aligned} P &= \frac{1}{32\Lambda^2 T} \int_0^\infty \frac{k^2 dk}{2\pi^2} \left[k^4 Q_{ij}(k) Q_{kl}(-k) \frac{2}{5} \left(\delta_{ik}\delta_{jl} - \frac{4}{3}\delta_{ij}\delta_{kl} \right) \right. \\ &\quad \left. + \frac{16}{9} k^4 J_{ij}(k) J_{kl}(-k) \frac{2}{5} \left(\delta_{ik}\delta_{jl} - \frac{4}{3}\delta_{ij}\delta_{kl} \right) \right. \\ &\quad \left. + \frac{k^6}{9} O_{ijk}(k) O_{lmn}(-k) \frac{2}{21} (\delta_{il}\delta_{jm}\delta_{kn} + \text{trace terms}) \right] \\ &= \frac{2G_N}{5} \int_0^\infty \frac{d\omega}{2\pi} \omega^6 \left[|I_{ij}(\omega)|^2 + \frac{16}{9} |J_{ij}(\omega)|^2 + \frac{5}{189} \omega^2 |I_{ijk}(\omega)|^2 + \dots \right] \end{aligned} \quad (202)$$

which, once averaged over time, recovers at the lowest order the standard Einstein quadrupole formula plus magnetic quadrupole and electric octupole contributions

$$P = \frac{G_N}{5} \langle \ddot{Q}_{ij}^2 \rangle + \frac{16G_N}{45} \langle \ddot{J}_{ij} \rangle + \frac{G_N}{189} \langle \ddot{O}_{ijk} \rangle \quad (203)$$

where we have used that


$$\int_0^\infty \frac{d\omega}{2\pi} |\tilde{A}(\omega)|^2 = \frac{1}{2} \int_{-\infty}^\infty dt A^2(t).$$

There are however corrections to this result for any given multipole, due to the scattering of the GW off the curved space-time because of the presence of the static potential due to the presence of the massive binary system. The first of such corrections scale as $G_N M \int (d^4 k \frac{1}{k})^2 \delta^3(k) \sim G_N M k \sim v^3$ (for radiation $k \sim v/r$), that is a 1.5PN correction with respect to the leading order. The tail amplitude is described by the diagram in fig. 14 and it adds up to the leading order to give a contribution to the flux going as

$$\left| \frac{A_h|_{v^3}}{A_h|_{v^0}} \right|^2 = 1 + 2\pi G_N M \omega + O(v^6). \quad (204)$$

Finally one could consider the scattering of the emitted GW wave off another GW, as in fig. 15. This process is known as *non-linear memory* effect, it represents a 2.5PN correction with respect to the leading emission amplitude⁴⁹ and it has not yet been computed within the effective field theory formalism.

Combined tail and memory effects enter at 4PN order in the emitted radiation, i.e. double scattering of the emitted radiation off the background curvature *and* off another GW. The divergences describing such process have been analyzed in⁵⁰, leading to the original derivation of the mass renormalization described in subsec. . The renormalization group equations allow a resummation of the logarithmic term making a non-trivial prediction for the pattern of the leading UV logarithms appearing at higher orders⁵¹.



⁴⁹ D. Christodoulou. Nonlinear nature of gravitation and gravitational wave experiments. *Phys.Rev.Lett.*, 67:1486–1489, 1991. DOI: 10.1103/PhysRevLett.67.1486; Luc Blanchet and Thibault Damour. Hereditary effects in gravitational radiation. *Phys.Rev.*, D46:4304–4319, 1992. DOI: 10.1103/PhysRevD.46.4304; and Luc Blanchet. Gravitational wave tails of tails. *Class.Quant.Grav.*, 15:113–141, 1998. DOI: 10.1088/0264-9381/15/1/009

⁵⁰ A. Ross W. D. Goldberger and I. Z. Rothstein. Black hole mass dynamics and renormalization group evolution. arXiv:1211.6095 [hep-th], 2012

⁵¹ W. D. Goldberger and A. Ross. Gravitational memory corrections from emitted from a source, scattered by another GW, before reaching the observer. *Phys.Rev.D*, 81:124015, 2010, and A. Ross W. D. Goldberger and I. Z. Rothstein. Black hole mass dynamics and renormalization group evolution. arXiv:1211.6095 [hep-th], 2012

Hints of data analysis

Using that for a binary system in circular orbit the relevant quadrupole moment components are

$$\begin{aligned} M_{11} &= \frac{\mu}{2} \omega_{GW}^2 R^2 \cos(\omega_{GW} t), \\ M_{12} &= \frac{\mu}{2} \omega_{GW}^2 R^2 \sin(\omega_{GW} t), \end{aligned} \quad (205)$$

where the reduced mass $\mu \equiv m_1 m_2 / (m_1 + m_2)$, we have now gathered all the elements to compute the GW-form, which is

$$\begin{aligned} h_+ &= \frac{1}{r} G_N \mu (2\pi R f_{GW})^2 F_+(t) \cos(\Phi(t)), \\ &= \frac{4}{r} \eta (G_N M)^{5/3} (\pi f_{GW})^{2/3} F_+(t) \cos(\Phi(t)) \end{aligned} \quad (206)$$

where the Kepler's law $(\omega_{GW}/2)^2 = G_N M / R^3$. Note that both $h(t)$ as increasing amplitude and instantaneous frequency. Its Fourier transform can be computed via the stationary phase approximation, see ex. 35 to give

$$\tilde{h}_+(f) = \left(\frac{5}{24\pi^4/3} \right)^{1/2} e^{i\psi(f)} \frac{1}{r} \eta^{1/2} \frac{(G_N M)^{5/6}}{f^{7/6}} F_+(t) \quad (207)$$

Given the noise characteristic of the detector, quantified by the noise spectral density $S_n(f)$ defined by the average over noise realiations

$$\langle n(f)n(f') \rangle \equiv S_n(f) \delta(f - f') \quad (208)$$

one can define the Signal-to-Noise Ratio (SNR) of a signal as

$$\begin{aligned} SNR^2 &= \int df \frac{|h(f)|^2}{S_n(f)} \\ &= \int d \ln f \frac{f |h(f)|^2}{S_n(f)} \end{aligned} \quad (209)$$

Exercise 27 ***** Multipole decomposition

Derive eq. (161).

Hint: use the trick

$$\int d^3x (T_{ij}x_k + T_{ik}x_j) = \int d^3x T_{il} (x_j x_k)_{,l}$$

to derive, via $\dot{T}_{0i} = -T_{il,l}$, that

$$\int d^3x (T_{ij}x_k + T_{ik}x_j) = \int d^3x \dot{T}_{i0} x_j x_k.$$

Now use the same trick to derive

$$\int d^3x \dot{T}_{i0} x_j x_k + \dot{T}_{j0} x_k x_i + \dot{T}_{k0} x_i x_j = \int d^3x \ddot{T}_{00} x_i x_j x_k.$$

Exercise 28 **** Multipole reduction

Derive eq. (161).

Hint: use a trick analog to eq. (150) to derive

$$\int d^3x \left[T_{ij}x_k + T_{ki}x_j + T_{jk}x_i \right] = \frac{1}{2} \int d^3x \left[\dot{T}_{i0}x_jx_k + \dot{T}_{k0}x_ix_j + \dot{T}_{k0}x_ix_j \right]$$

and then use

$$\begin{aligned} \int d^3x \left[T_{ij}x_k - T_{ik}x_j \right] &= \int d^3x \left[T_{il}x_{j,l}x_k - T_{kl}x_{i,l}x_j \right] \\ &= \int d^3x \left[\dot{T}_{i0}x_jx_k - T_{ik}x_j - \dot{T}_{k0}x_ix_j + T_{kj}x_i \right] \\ \implies \int d^3x \left[T_{ij}x_k - T_{jk}x_i \right] &= \int d^3x \left[\dot{T}_{i0}x_jx_k - \dot{T}_{k0}x_ix_j \right] \\ \implies \int d^3x \left[T_{ij}x_k - T_{ki}x_j \right] &= \int d^3x \left[\dot{T}_{j0}x_ix_k - \dot{T}_{k0}x_ix_j \right] \end{aligned}$$

Now combine the 3 equations involving $T_{ij}x_k$ to derive the result.

Exercise 29 Velocity quadrupole coupling to the Weyl tensor

From eq. 158 derive the formulae

$$\begin{aligned} E_{ij} &= \frac{1}{2} \ddot{\sigma}_{ij} \\ B_{ij} &= \frac{1}{4} \epsilon_{ikl} \left(\dot{\sigma}_{kj,l} - \dot{\sigma}_{lj,k} \right) \end{aligned}$$

valid at linear order in the TT gauge. From the expression of B_{ij} derive eq. (162).

Hint: express the contraction of the second line of eq. (161) with $\sigma_{ij,k}$ as

$$\begin{aligned} & \left(\dot{T}_{0i}x_jx_k + \dot{T}_{0j}x_ix_k - 2\dot{T}_{0k}x_ix_j \right) = \\ & T_{0i}x_jx_k \left(\delta^{im} \delta^{kn} - \delta^{in} \delta^{km} \right) + T_{0j}x_ix_k \left(\delta^{jm} \delta^{kn} - \delta^{jn} \delta^{km} \right). \end{aligned}$$

Now substitute $\delta^{ac} \delta^{bd} - \delta^{ad} \delta^{bc} = \epsilon^{abi} \epsilon^{cdi}$ to obtain the last term in eq. (162).

Exercise 30 ***** Alternative derivation of the Burke-Thorne potential

Go through eq.(176) and contract the two Weyl tensor using their expression in the TT gauge:

$$R_{i0j0}^0 = \frac{1}{2} \ddot{\sigma}_{ij}^{(TT)}.$$

Hint: Remember that the propagator (132) for σ is not the same as the one for $\sigma_{(TT)}$, as

$$\sigma_{ij}^{(TT)} = \Lambda_{ij,kl} \sigma_{kl},$$

with $\Lambda_{ij,kl}$ given by eq.(22).


Exercise 31 ***** Time dependence of total mass of a binary system

In order to compute the mass of a binary system in terms of its binary constituents properties check how the h_{00} components of the gravitational field couple to it. At leading order the diagram we have the first diagram in fig. 16 where the field ϕ couple to the monopole mass. Neglecting the angular momentum (spin) of the composite object, the next diagram comes from the second and third graphs in fig. 16. For computing the last diagram it will be useful the expansion of R^0_{i0j} at linear order in ϕ and σ , i.e.

$$\begin{aligned} R^0_{i0j} = & \frac{1}{2}\ddot{\sigma}_{ij} - \delta_{ij}\ddot{\phi} - \phi_{ij} - \frac{1}{2}A_{i,j} - \frac{1}{2}A_{j,i} \\ & + \frac{1}{2}\phi_k(\sigma_{ik,j} + \sigma_{jk,i} - \sigma_{ij,k}) - \frac{3}{2}\dot{\phi}\dot{\sigma}_{ij} - 2\phi\ddot{\sigma}_{ij} - \ddot{\phi}\sigma_{ij} \\ & + \delta_{ij}(A_k\dot{\phi}_k + \phi_k\dot{A}_k) - 2A_j\dot{\phi}_i - 2A_i\dot{\phi}_j - 2\phi_i\dot{A}_j - 2\phi_j\dot{A}_i - \frac{1}{2}(A_{i,j} + A_{j,i})\dot{\phi} \\ & + \delta_{ij}\phi_k\phi_k + 2\delta_{ij}\phi^2 - 3\phi_i\phi_j + 4\phi\dot{\phi} + O(h^3) \end{aligned}$$

and using the ‘‘bulk’’ vertices of ex. 25.

Result: see eq.(8) of ⁵².



⁵² A. Ross W. D. Goldberger and I. Z. Rothstein. Black hole mass dynamics and renormalization group evolution. Figure 16: Coupling of gravity to a binary system at leading order and at Q^2 order. arXiv:1211.6009 [hep-th] 2012

Exercise 32 *** Scaling of GW emission diagrams

Derive the scaling of the diagram in fig. 13.

Result: $d\omega d^3k \omega^2 \tilde{Q}(\omega) H(\omega, \mathbf{k})$.

Derive the scaling of the diagram in fig. 14. Hint: use

- the scaling of the vertex at the GW emission above (divided by Λ)
- the scaling of the GW propagator $\delta^{(4)}(k)/k^2$
- the scaling of the Newtonian graviton emission $dtd^3q M\phi(t, \mathbf{q})/\Lambda$
- the scaling of the ϕ propagator $\delta(t)/q^2$
- the scaling of the $\sigma^2\phi$ vertex: $dtd^3x H(t, \mathbf{x}) \int d\omega d^3k d^3q \omega^2$.

Result: the amplitude scales as $G_N H(\omega, \mathbf{k}) M \tilde{Q}(\omega) d\omega d^3k \omega^3$, i.e. as $G_N M \omega \sim v^3$ times the above one.

Exercise 33 **** Memory effect

The amplitude for emission of GWs receives corrections from the diagram in fig. 15, where GWs are emitted from two different quadrupole-gravity vertices and then interact together. Adapt the scaling rules described in ex. 32 to this case, in which circulating momenta have $k_0 \sim k \sim v/r$ and the sources emitting the GW $\ddot{Q} \sim Mv^2 \sim d\omega\omega^2\tilde{Q}(\omega)$.

Result: $G_N H(\omega, \mathbf{k}) d\omega d^3k \omega^5 \tilde{Q}(\omega) d\omega \tilde{Q}(\omega)$, i.e. of order v^5 with respect leading order diagram in fig. 13.

Exercise 34 ** Energy from propability amplitude

Check that eq. (198) has the correct dimensions

Exercise 35 ***** Fourier transform of the GW-form

From eq. (206) perform the Fourier integral

$$\tilde{h}_+(f) = \int df h_+(t) e^{2\pi i f t}$$

to obtain eq. (207).

Hint: the integral can be done using the stationary phase approximation, i.e.

$$\begin{aligned} \tilde{h}_+(f) &\propto \frac{1}{2} \int e^{2\pi i f t - i\Phi(t)} dt \\ &\simeq \frac{1}{2} e^{2\pi i f t_* - i\Phi(t_*)} \int e^{-i\ddot{\Phi}(t_*)(t-t_*)^2/2} dt \\ &= \frac{1}{2} e^{2\pi i f t_* - i\Phi(t_*)} \left(\frac{2\pi}{\ddot{\Phi}(t_*)} \right) e^{-i\pi/4} \end{aligned}$$

where t_* is the solution of $2\pi f = \dot{\Phi}(t_*)$ and in the last passage the complex Gaussian integral has been performed. Substituting the expression for the phase as a function of $t_*(f)$ as it can be inferred from eq. (99), one obtains eq. (207).

Exercise 36 ***** Polarization sum

For the magnetic quadrupole, the polarization tensor sum boils down to:

$$\begin{aligned} &\int \frac{d\Omega}{4\pi} \sum_h \epsilon^{imn} n_n \epsilon^{kpq} n_q \epsilon_{ij}^*(\mathbf{k}, h) \epsilon_{kl}(\mathbf{k}, h) \\ &= \int \frac{d\Omega}{4\pi} \frac{n_n n_q}{2} \left[(\delta_{mp} \delta_{jl} + \delta_{ml} \delta_{jp} - \delta_{mj} \delta_{pl}) + n_j n_l n_m n_p \right. \\ &\quad \left. - (\delta_{jl} n_m n_p + \delta_{mp} n_j n_l + \delta_{ml} n_j n_p + \delta_{jp} n_m n_l - \delta_{mj} n_p n_l - \delta_{pl} n_m n_j) \right] \quad (210) \\ &= \frac{1}{5} \left(\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk} - \frac{2}{3} \delta_{ij} \delta_{kl} \right). \end{aligned}$$

For the octupole the analog sum is given by (symmetrization under $i \leftrightarrow j$ and $l \leftrightarrow m$ is understood, however since it is going to multiply tensors which are symmetric under such exchange, we do

not need to worry about that):

$$\begin{aligned}
 & \int d\Omega 4\pi \sum_h n^k n^n \epsilon_{ij}^*(\mathbf{k}, h) \epsilon_{lm}(\mathbf{k}, h) \\
 &= \int d\Omega n_m n_n \left(\delta_{il} \delta_{jm} - n_i n_l \delta_{jm} - n_j n_m \delta_{il} + \frac{1}{2} n_i n_j n_l n_m - \frac{1}{2} \delta_{ij} \delta_{lm} + \frac{1}{2} (n_i n_j \delta_{lm} + n_l n_m \delta_{ij}) \right) \quad (211) \\
 &= \frac{2}{21} \left[\delta_{il} \delta_{jm} \delta_{kn} - \frac{1}{5} (\delta_{il} \delta_{jk} \delta_{mn} + \delta_{im} \delta_{ln} \delta_{jk} + \delta_{in} \delta_{lm} \delta_{jk}) \right]
 \end{aligned}$$

Derive from the sums above the relative term in eq. (202).

Exercise 37 * Derivation of the flux formula in terms of binary velocity)**

Use the leading quadrupole term in eq.(202) to derive the leading order of the flux formula eq. (3) for circular orbits.

Hint: use eq. (205) and that

$$v = (G_N M \pi f_{GW})^{1/3}.$$

List of Figures

- 1 Interferometer scheme. Light emitted from the laser is shared by the two orthogonal arms after going through the beam splitter. After bouncing at the end mirrors it is recombined at the photo-detector. 21
- 2 Feynman graph accounting for the Newtonian potential. 33
- 3 Diagram giving a power-law divergent contribution to the mass. 34
- 4 Graph giving a G_N^2 contribution to the 1PN potential via ϕ propagators. 36
- 5 Vertex scaling: $\frac{m}{\Lambda} dt d^d k \sim dt \frac{m}{\Lambda} r^{-d}$ 37
- 6 A Green function is represented by a propagator, with scaling: $\delta(t) \delta^d(k) / k^2 \sim \delta(t) r^{2+d}$ 37
- 7 Triple internal vertex scaling: $\frac{(k^2, k k_0, k_0^2)}{\Lambda} \delta^d(k) dt (d^d k)^3 \sim dt \frac{(1, v, v^2)}{r^{2+2d} \Lambda}$ 37
- 8 Quantum contribution to the 2-body potential. 38
- 9 Sub-leading correction to the 2 particle scattering process due to gravity self-interaction. The ϕ^3 vertex brings two time derivative, making this diagram contribute from 2PN order on. 40
- 10 Graph dressing T_{00} at v^2 order and T_{ij} at leading order. The external radiation graviton does not carry momentum but it is Taylor expanded. 50
- 11 Diagram giving the leading term of the amplitude describing radiation back-reaction on the sources. 50
- 12 Next-to-leading order term in the back-reaction amplitude. 51
- 13 Diagram representing the emission of a GW from a quadrupole source 57
- 14 Emission of a GW from a quadrupole source with post-Minkowskian correction represented by the scattering off the background curved by the presence of binary system. 58
- 15 Memory diagram: GW emitted from a source scattered by another GW before reaching the observer. 60
- 16 Coupling of gravity to a binary system at leading order and at Q^2 order. 63

List of Tables

Bibliography

T. Appelquist and J. Carazzone. Infrared singularities and massive fields. *Phys. Rev. D*, 11:2856, 1975.

L. Blanchet. Time asymmetric structure of gravitational radiation. *Phys.Rev.*, D47:4392–4420, 1993.

L. Blanchet and Gerhard Schafer. Gravitational wave tails and binary star systems. *Class.Quant.Grav.*, 10:2699–2721, 1993.

L. Blanchet, S. L. Detweiler, A. Le Tiec, and B. F. Whiting. High-order post-newtonian fit of the gravitational self-force for circular orbits in the schwarzschild geometry. *Phys. Rev.*, D81, 2010.

Luc Blanchet. Gravitational wave tails of tails. *Class.Quant.Grav.*, 15: 113–141, 1998. DOI: 10.1088/0264-9381/15/1/009.

Luc Blanchet. Gravitational radiation from post-newtonian sources and inspiralling compact binaries. *Living Reviews in Relativity*, 9(4), 2006. URL <http://www.livingreviews.org/lrr-2006-4>.

Luc Blanchet and Thibault Damour. Tail transported temporal correlations in the dynamics of a gravitating system. *Phys.Rev.*, D37: 1410, 1988.

Luc Blanchet and Thibault Damour. Hereditary effects in gravitational radiation. *Phys.Rev.*, D46:4304–4319, 1992. DOI: 10.1103/Phys-RevD.46.4304.

Alessandra Buonanno. Gravitational waves. 2007. URL <http://arxiv.org/abs/arXiv:0709.4682>.

M. Burgay, N. D’Amico, A. Possenti, R. N. Manchester, A. G. Lyne, B. C. Joshi, M. A. McLaughlin, and M. Kramer *et al.* An increased estimate of the merger rate of double neutron stars from observations of a highly relativistic system. *Nature*, 426:531, 2003.

W. L. Burke and K. S. Thorne. In S. I. Fickler M. Carmeli and L. Witten, editors, *Relativity*, pages 209–228. Plenum, New York, 1970.

M. Tiglio C. R. Galley. Radiation reaction and gravitational waves in the effective field theory approach. *Phys. Rev.*, D79:124027, 2009.

D. Christodoulou. Nonlinear nature of gravitation and gravitational wave experiments. *Phys.Rev.Lett.*, 67:1486–1489, 1991. DOI: 10.1103/PhysRevLett.67.1486.

T. Damour. Gravitational radiation and the motion of compact bodies. In N. Deruelle and T. Piran, editors, *Gravitational Radiation*, pages 59–144. North-Holland, Amsterdam, 1983a.

T. Damour. *Gravitational radiation and the motion of compact bodies*, pages 59–144. North-Holland, Amsterdam, 1983b.

B. DeWitt. Effectice action for expectation values. In R. Penrose and C. J. Isham, editors, *Quantum concepts in Space and Time*. Clarendon Press, Oxford, 1986.

Eanna E. Flanagan and Scott A. Hughes. The basics of gravitational wave theory. *New J. Phys.*, 7:204, 2005.

S. Foffa and R. Sturani. Tail terms in gravitational radiation reaction via effective field theory. *Phys. Rev. D*, 87:044056, 2013.

Stefano Foffa and Riccardo Sturani. Effective field theory calculation of conservative binary dynamics at third post-Newtonian order. *Phys.Rev.*, D84:044031, 2011.

Stefano Foffa and Riccardo Sturani. Effective field theory methods to model compact binaries. *Class.Quant.Grav.*, 31(4):043001, 2014. DOI: 10.1088/0264-9381/31/4/043001.

Andreas Freise and Kenneth A. Strain. Interferometer techniques for gravitational-wave detection. *Living Reviews in Relativity*, 13(1), 2010. DOI: 10.12942/lrr-2010-1. URL <http://www.livingreviews.org/lrr-2010-1>.

C. R. Galley. The classical mechanics of non-conservative systems. *Phys. Rev. Lett.*, 110:174301, 2013.

C. R. Galley and A. K. Leibovich. Radiation reaction at 3.5 post-newtonian order in effective field theory. *Phys. Rev. D*, 86:044029, 2012.

W. D. Goldberger. Les houches lectures on effective field theories and gravitational radiation. In *Les Houches Summer School - Session 86: Particle Physics and Cosmology: The Fabric of Spacetime*, 2007.

W. D. Goldberger and A. Ross. Gravitational radiative corrections from effective field theory. *Phys. Rev. D*, 81:124015, 2010.

- Walter D. Goldberger and Ira Z. Rothstein. An Effective field theory of gravity for extended objects. *Phys.Rev., D73*:104029, 2006.
- J.J. Hermes, Mukremin Kilic, Warren R. Brown, D.E. Winget, Carlos Allende Prieto, et al. Rapid Orbital Decay in the 12.75-minute WD+WD Binary J0651+2844. 2012. DOI: 10.1088/2041-8205/757/2/L21.
- R A Hulse and J H Taylor. Discovery of a pulsar in a binary system. *Astrophys. J.*, 195:L51, 1975.
- C. Itzinkinson and J. B. Zuber. *Quantum field theory*. Mac Graw-Hill International Book Company, 1980.
- B. R. Iyer and C. M. Will. Postnewtonian gravitational radiation reaction for two-body systems. *Phys. Rev. Lett.*, 70:113, 1993.
- John David Jackson. *Classical Electrodynamics*. John Wiley & Sons, iii edition, 1999.
- L. V. Keldysh. Diagram technique for nonequilibrium processes. *Zh. Eksp. Teor. Fiz.*, 47:1515–1527, 1964.
- Barak Kol and Michael Smolkin. Black hole stereotyping: Induced gravito-static polarization. *JHEP*, 1202:010, 2012. DOI: 10.1007/JHEP02(2012)010.
- M. Kramer and N. Wex. The double pulsar system: A unique laboratory for gravity. *Class. Quant. Grav.*, 26:073001, 2009.
- M. Maggiore. *Gravitational Waves*. Oxford University Press, 2008.
- A. Ross R. A. Porto and I. Z. Rothstein. Spin induced multipole moments for the gravitational wave amplitude from binary inspirals to 2.5 post-newtonian order. *JCAP*, 1209:028, 2012.
- B.S. Sathyaprakash and B.F. Schutz. Physics, Astrophysics and Cosmology with Gravitational Waves. *Living Rev.Rel.*, 12:2, 2009.
- J. S. Schwinger. Brownian motion of a quantum oscillator. *J. Math. Phys.*, 2:407–432, 1961.
- LIGO Scientific and Virgo Collaborations. Predictions for the rates of compact binary coalescences observable by ground-based gravitational-wave detectors. *Class. Quant. Grav.*, 27:173001, 2010.
- I. H. Stairs, S. E. Thorsett, J. H. Taylor, and A. Wolszczan. Studies of the relativistic binary pulsar psr b1534+12: I. timing analysis. *Astrophys. J.*, 581:501, 2002.

R Sturani. Ligo/virgo/geo/kagra science. In P Binétruy, G Auger, and E. Plagnol, editors, 9th *LISA Symposium, Paris*, volume 467 of *ASP Conference Series*, pages 293–302. Astronomical Society of the Pacific, 2012. URL <http://www.apc.univ-paris7.fr/Downloads/lisa/aspvolume.pdf>.

A. Ross W. D. Goldberger and I. Z. Rothstein. Black hole mass dynamics and renormalization group evolution. arXiv:1211.6095 [hep-th], 2012.

J. M. Weisberg and J. H. Taylor. Observations of post-newtonian timing effects in the binary pulsar psr 1913+16. *Phys. Rev. Lett.*, 52: 1348, 1984.

A. Wolszczan. A nearby 37.9-ms radio pulsar in a relativistic binary system. *Nature*, 350:688, 1991.