TECHNISCHE UNIVERSITÄT DARMSTADT

Nuclear Reactions and Multineutron Correlations

Hans-Werner Hammer
Technische Universität Darmstadt

HFHF

Bundesministerium für Bildung und Forschung

DFG Deutsche Forschungsgemeinschaft

SFB 1245

Workshop on Multineutron Clusters in Nuclei and in Stars, ICTP-SAIFR, Sao Paulo, June 2-6, 2025

Outline

TECHNISCHE UNIVERSITÄT DARMSTADT

- Universality and the unitary limit
- Schrödinger symmetry
- Nuclear reactions with neutrons
- Summary and Outlook
- (Neutral charm mesons and the $X(3872)$)

References:

HWH, D.T. Son, Proc. Nat. Acad. Sci. **118**, e2108716118 (2021)
T. Backert, S. Dietz, HWH, S. König, D.T. Son, in preparation
M. Göbel, HWH, D.R. Phillips, Phys. Rev. C **110**, 024003 (2024)

Universality

TECHNISCHE UNIVERSITÄT DARMSTADT

Universality: Physical systems with different short-distance behavior exhibit identical behavior at large distances

Universality

Universality: Physical systems with different short-distance behavior exhibit identical behavior at large distances

- Condensed matter systems near critical point



$$\rho_{liq/gas}(T) - \rho_c \longrightarrow \pm A(T_c - T)^\beta$$

Liquid-gas system



$$M_0(T) \longrightarrow A'(T_c - T)^\beta$$

Ferromagnet (one easy axis)

$\beta = 0.325$

- Universality class determines critical exponents:
- Scale invariance (often conformal invariance)

Physics Near the Unitary Limit

TECHNISCHE UNIVERSITÄT DARMSTADT

- Consider short-ranged, resonant S-wave interactions $\Longrightarrow$ halo state
- Unitary limit: $a \to \infty, \ell \sim r_e \to 0$

$$\mathcal{T}_2(k,k) \propto \Big[\underbrace{k \cot \delta}_{-1/a + r_e k^2/2 + \ldots} - ik \Big]^{-1} \sim i/k$$





- Scattering amplitude scale invariant, saturates unitarity bound

Physics Near the Unitary Limit

TECHNISCHE UNIVERSITÄT DARMSTADT

- Consider short-ranged, resonant *S*-wave interactions $\Longrightarrow$ halo state
- Unitary limit: $a \to \infty, \ell \sim r_e \to 0$

$$\mathcal{T}_2(k,k) \propto \Big[\underbrace{k \cot\delta}_{-1/a + r_e k^2/2 + \ldots} - ik \Big]^{-1} \sim i/k$$





  - Scattering amplitude scale invariant, saturates unitarity bound
- BEC/BCS crossover, neutron matter, ... (cf. Schäfer, Baym, PNAS **118**, e2113775118 (2021))



BCS BEC Crossover - Randeria, Taylor, 2014



Sebastian König



Robert Schulze CC BY-SA 3.0, via Wikimedia Commons

Physics Near the Unitary Limit

TECHNISCHE UNIVERSITÄT DARMSTADT

- Consider short-ranged, resonant S-wave interactions $\Longrightarrow$ halo state
- Unitary limit: $a \to \infty, \ell \sim r_e \to 0$

$$\mathcal{T}_2(k,k) \propto \Big[\underbrace{k \cot\delta}_{-1/a + r_e k^2/2 + \ldots} - ik \Big]^{-1} \sim i/k$$





  - Scattering amplitude scale invariant, saturates unitarity bound

- System has also (non-relativistic) conformal symmetry/Schrödinger symmetry
  Mehen, Stewart, Wise, PLB **474**, 145 (2000); Nishida, Son, PRD **76**, 086004 (2007); ...
- Exploit approximate conformal symmetry for nuclear reactions with neutrons

$$a \approx -18.6 \text{ fm}, \quad r_e \approx 2.8 \text{ fm} \quad \Rightarrow \quad \underbrace{1/(ma^2)}_{0.1 \text{ MeV}} \ll E_n^{cms} \ll \underbrace{1/(mr_e^2)}_{5 \text{ MeV}}$$

Schrödinger symmetry

TECHNISCHE UNIVERSITÄT DARMSTADT

- Non-relativistic conformal symmetry: Schrödinger symmetry
  - Galilei symmetry
    - space + time translations
    - rotations
    - Galilei boosts
  - Scale transformations

$$\boldsymbol{x} \to e^{\lambda}\boldsymbol{x}, \qquad t \to e^{2\lambda}t, \qquad \psi \to e^{-\lambda\Delta}\psi$$

  - Special conformal transformations

$$\boldsymbol{x} \to \frac{\boldsymbol{x}}{1+\xi t}\,, \qquad t \to \frac{t}{1+\xi t}\,, \qquad \psi \to \psi' = \ldots$$

  $\Rightarrow$ preserves angles
- 12 Parameters
- Generators: $H, \boldsymbol{P}, \boldsymbol{L}, \boldsymbol{K}, D, C,$ satisfy Schrödinger algebra

Unitary limit: RG perspective

TECHNISCHE UNIVERSITÄT DARMSTADT

- Spin-1/2 Fermions with zero-range interactions ($|a| \gg r_e \sim \ell$)



- Renormalization group equation: $\Lambda \dfrac{d}{d\Lambda} \tilde{g}_2 = \tilde{g}_2(1 + \tilde{g}_2)$
- Two fixed points:
  - $\tilde{g}_2 = 0 \;\Leftrightarrow\; a = 0 \quad \Rightarrow$ no interaction, free particles
  - $\tilde{g}_2 = -1 \;\Leftrightarrow\; 1/a = 0 \quad \Rightarrow$ unitary limit $\quad \Rightarrow$ conformal/Schrödinger symmetry
- Neutrons: $a \approx -18.6$ fm, $\quad r_e \approx 2.8$ fm $\quad \Rightarrow$ close to unitary limit $\quad \Rightarrow$ approximately conformal

$\Longrightarrow$ multineutron system can be described by conformal field

Conformal field theory constraints

TECHNISCHE UNIVERSITÄT DARMSTADT

- Two-point function of (primary) conformal field operator $\mathcal{U}$

$$G_{\mathcal{U}}(t,\boldsymbol{x}) = -i\langle T\mathcal{U}(t,\boldsymbol{x})\mathcal{U}^\dagger(0,\boldsymbol{0})\rangle = C\,\frac{\theta(t)}{(it)^\Delta}\exp\left(\frac{iM\boldsymbol{x}^2}{2t}\right)$$

- Determined by symmetry up to overall constant C
- Two-point function in momentum space

$$G_{\mathcal{U}}(\omega,\boldsymbol{p}) = -C\left(\frac{2\pi}{M}\right)^{3/2}\Gamma\left(\frac{5}{2}-\Delta\right)\left(\frac{\boldsymbol{p}^2}{2M}-\omega\right)^{\Delta-\frac{5}{2}}$$

  - pole only for $\Delta = 3/2$ (free field)
  - branch cut for $\Delta > 3/2$

Conformal field theory constraints

TECHNISCHE UNIVERSITÄT DARMSTADT

- Imaginary part of propagator

$$\mathrm{Im}\, G_{\mathcal{U}}(\omega, \boldsymbol{p}) \sim \begin{cases} \delta\left(\omega - \frac{\boldsymbol{p}^2}{2M}\right), & \Delta = \frac{3}{2}\,, \\ \left(\omega - \frac{\boldsymbol{p}^2}{2M}\right)^{\Delta - \frac{5}{2}} \theta\left(\omega - \frac{\boldsymbol{p}^2}{2M}\right), & \Delta > \frac{3}{2} \end{cases}$$

- General $\mathcal{U}$ does not behave like a particle ("unparticle/unnucleus") $\Rightarrow$ continuous energy spectrum
- Examples of unnuclei
  - N free fields: $\mathcal{U} = \psi_1 \ldots \psi_N, \quad M = N m_\psi, \quad \Delta = 3N/2$
  - N interacting fields: $\mathcal{U} = \psi_1 \ldots \psi_N, \quad M = N m_\psi, \quad \Delta > 3/2$
- Corrections from finite a and r_0 (S. Dutta, R. Mishra, D.T. Son, Phys. Rev. D **109**, 016001 (2024))

$$\mathrm{Im}\, G_{\mathcal{U}}(\omega, 0) \sim \omega^{\Delta - \frac{5}{2}} \theta(\omega) \left(1 + \frac{c_1}{a\sqrt{m\omega}} + c_2 r_0 \sqrt{m\omega}\right), \quad c_2 = 0$$

Scaling dimension $\Leftrightarrow$ critical exponents

TECHNISCHE UNIVERSITÄT DARMSTADT

- How to calculate scaling dimension Δ? not given by symmetry $\Longrightarrow$ non-perturbative problem
  - (1) Δ can be obtained from field theory calculation
  - (2) Δ can be obtained from operator state correspondence

Δ of primary operator = (Energy of state in HO)/$\hbar\omega$

(Nishida, Son, Phys. Rev. D **76**, 086004 (2007))

N	S	L	$\mathcal{O}$	Δ	Δ_{free}
2	0	0	$\psi_1\psi_2$	2	3
3	1/2	1	$\psi_1\psi_2\nabla_j\psi_2$	4.27272	5.5
3	1/2	0	$\psi_1\nabla_j\psi_2\nabla_j\psi_2$	4.66622	6.5
4	0	0	$\psi_1\psi_2\nabla_j\psi_1\nabla_j\psi_2$	5.07(1)	8
5	1/2	1	$\psi_1\psi_2\nabla_j\psi_1\nabla_j\psi_2\nabla_i\psi_1$	7.6(1)	10.5

- Connection between Δ and energy of particles in a trap
- Predictions for $N \leq 70$ available (M.G. Endres et al., PRA **84**, 043644 (2011))

Reactions with neutrons

TECHNISCHE UNIVERSITÄT DARMSTADT

- High-energy nuclear reaction with multi-neutron final state
  (HWH, Son, Proc. Nat. Acad. Sci. **118**, e2108716118 (2021))



$$\Delta Mc^2 + E_{A_1} + E_{A_2} = E_B + E_{\mathcal{U}}$$

- Assumption: energy scale of primary reaction $\gg E_{\mathcal{U}} - \frac{\boldsymbol{p}^2}{2M_{\mathcal{U}}} = E_n^{cms}$
- Factorization: $\dfrac{d\sigma}{dE} \sim |\mathcal{M}_{primary}|^2 \,\mathrm{Im}\, G_{\mathcal{U}}(E_{\mathcal{U}}, \boldsymbol{p})$
- Reproduces Watson-Migdal treatment of FSI for $2n$

Conformal symmetry and reactions with neutrons

TECHNISCHE UNIVERSITÄT DARMSTADT

- Two ways to do experiments
  - (a) detect recoil particle B $\quad \frac{d\sigma}{dE} \sim (E_0 - E_B)^{\Delta - 5/2}, \qquad E_0 = (1 + M_B/M_{\mathcal{U}})^{-1} E_{\rm kin}$
  - (b) detect all final state particles including neutrons: $\quad \frac{d\sigma}{dE} \sim (E_{xn}^{cms})^{\Delta - 5/2}$

Conformal symmetry and reactions with neutrons

TECHNISCHE UNIVERSITÄT DARMSTADT

- Two ways to do experiments
  - (a) detect recoil particle B $\quad \dfrac{d\sigma}{dE} \sim (E_0 - E_B)^{\Delta-5/2}, \qquad E_0 = (1 + M_B/M_{\mathcal{U}})^{-1} E_{\rm kin}$
  - (b) detect all final state particles including neutrons: $\quad \dfrac{d\sigma}{dE} \sim (E_{xn}^{cms})^{\Delta-5/2}$
- $2n$ case can be understood from dimer propagator in halo/pionless EFT ($\Delta = 2$)

$$G_d(E_{nn}, \mathbf{0}) \sim \frac{1}{1/a + i\sqrt{mE_{nn}}} \quad \Rightarrow \quad \mathrm{Im}\, G_d(E_{nn}, \mathbf{0}) \sim \frac{\sqrt{E_{nn}}}{(ma^2)^{-1} + E_{nn}}$$

- See also (M.Göbel et al., Phys. Rev. C **104**, 024001 (2021)) for $^6\mathrm{He}(p, p\alpha)2n$

Predictions for relative energy distributions

TECHNISCHE UNIVERSITÄT DARMSTADT

- Power law behavior at low energies



- $N = 5$: $E^8 \longrightarrow E^{5.1}$
- $N = 6$: $E^{10.5} \longrightarrow E^{6.2}$
- ...
- Predictions for $N \leq 70$ available
  (M.G. Endres et al., PRA **84**, 043644 (2011))
- No peak except for $2n$

Braaten, HWH, Phys. Rev. D **107**, 034017 (2023)

3*n* **reaction calculations**

- Radiative muon/pion capture on the triton (AV18 + UIX)



Golak et al., PRC **98**, 054001 (2018)



Golak et al., PRC **94**, 054001 (2016)

- Conformal prediction: $d\Gamma/dE \sim (E_{3n})^{4.27272-5/2} \sim (E_{3n})^{1.77272}, \qquad 0.1\,\text{MeV} \ll E_{3n} \ll 5\,\text{MeV}$

3*n* **system in pionless EFT**

TECHNISCHE UNIVERSITÄT DARMSTADT

- Point-production amplitude in pionless EFT



- Reproduces conformal behavior (Backert, Dietz, HWH, König, Son, preliminary) → T. Backert (Friday)

P-Wave





- Free and unitary limit exponents reproduced

3*n* distributions from experiment

TECHNISCHE UNIVERSITÄT DARMSTADT

- 3*n* case consistent with previous experiments for $^3\mathrm{H}(\pi^-,\gamma)3n$ (Miller et al., Nucl. Phys. A **343**, 347 (1980))
- 3*n* distributions from $^3\mathrm{H}(^3\mathrm{H},^3\mathrm{He})3n$ (Miki et al., Phys. Rev. Lett. **133**, 012501 (2024))





- Small q_{cm} data consistent

$4n$ distribution from experiment

- Search for tetraneutron resonances in $^{8}\mathrm{He}(p, p\alpha)4n$ (M. Duer et al., Nature **606**,678 (2022))





- Structure from initial state?
- Dineutron correlations can produce peak (Lazauskas, Hiyama, Carbonell, Phys. Rev. Lett. **130**, 102501 (2023))

4*n* distribution from experiment

TECHNISCHE UNIVERSITÄT DARMSTADT

- Search for tetraneutron resonances in $^{8}\mathrm{He}(p, p\alpha)4n$ (M. Duer et al., Nature **606**,678 (2022))





- Structure from initial state?
- Dineutron correlations can produce peak (Lazauskas, Hiyama, Carbonell, Phys. Rev. Lett. **130**, 102501 (2023))
- CFT result confirmed in pionless EFT (T. Backert, HWH, S. König, in preparation)

Initial state effects hard knock-out reactions: $2n$

TECHNISCHE UNIVERSITÄT DARMSTADT

- Universality of $2n$ distributions in hard core knock-out (Göbel, HWH, Phillips, Phys. Rev. C **110**, 024003 (2024))
- Calculate $2n$ distributions of different halo nuclei in halo EFT

	A	S_{2n} [keV]	E^*_{nc} [keV]	E^*_{nn} [keV]
^{11}Li	9	369	26	
^{14}Be	12	1266	510	
^{17}B	15	1384	83	118
^{19}B	17	90	5	
^{22}C	20	100	68	



- $S_{2n} \gg E^*_{nc}, E^*_{nn} \implies$ nuclei close to unitary limit $\implies$ shape accurate to 20%
- Initial state effects in conformal limit (D. Kromm)

Summary and Outlook

TECHNISCHE UNIVERSITÄT DARMSTADT

- Universality in strongly interacting quantum systems
- High-energy nuclear reactions with final state neutrons
  - $\Rightarrow$ (approximate) conformal symmetry
  - $\Rightarrow$ power law behavior of observables determined by scaling dimension Δ
- Model-independent constraints on (certain) nuclear reactions
- Connection between reactions & properties of trapped particles

Summary and Outlook

TECHNISCHE UNIVERSITÄT DARMSTADT

- Universality in strongly interacting quantum systems
- High-energy nuclear reactions with final state neutrons
  - ⇒ (approximate) conformal symmetry
  - ⇒ power law behavior of observables determined by scaling dimension Δ
- Model-independent constraints on (certain) nuclear reactions
- Connection between reactions & properties of trapped particles
- Understand difference between $3n$ and $4n$, initial state dependence
- Other applications & extensions
  - Neutral charm mesons
  - Two-component fermions in ultracold atom physics
  - Systems with the Efimov effect? (J. Rimek)
    - ⇒ bosonic atoms, nucleons, α particles
    - ⇒ only discrete scale symmetry, complex scaling dimensions

Additional transparencies

Unparticle physics

TECHNISCHE UNIVERSITÄT DARMSTADT

- (Relativistic) unparticle (Georgi, Phys. Rev. Lett. **98**, 221601 (2007))
  - field ψ in relativistic conformal field theory
  - ψ characterized by scaling dimension Δ, massless
  - hidden conformal symmetry sector beyond Standard model (weakly coupled)
  - no evidence at LHC so far
    (CMS Coll., EPJC **75**, 235 (2015), PRD **93**, 052011, JHEP **03**, 061 (2017))
- (Non-relativistic) unparticle/unnucleus
  - non-relativistic analog of Georgi's unparticle
  - field ψ in non-relativistic conformal field theory
    (cf. Nishida, Son, Phys. Rev. D **76**, 086004 (2007))
  - ψ characterized by scaling dimension Δ and mass M
  - free field has $\Delta = 3/2 \quad \Longleftrightarrow \quad$ mass dimension

    $\Rightarrow$ lowest possible value (unitarity)
  - N neutrons are (approximate) unparticle with mass Nm_N and scaling dimension $\Delta =$?

Hard knockout reaction calculations

TECHNISCHE UNIVERSITÄT DARMSTADT

- Two-neutron spectrum for $^6\mathrm{He}(p, p\alpha)2n$ (Göbel et al., Phys. Rev. C **104**, 024001 (2021))



- Can be understood from dimer propagator ($\Delta = 2$)

$$G_d(E_{nn}, \mathbf{0}) \sim \frac{1}{1/a + i\sqrt{mE_{nn}}} \quad \Rightarrow \quad \mathrm{Im}\, G_d(E_{nn}, \mathbf{0}) \sim \frac{\sqrt{E_{nn}}}{(ma^2)^{-1} + E_{nn}}$$

Neutral charm mesons and $X(3872)$

TECHNISCHE UNIVERSITÄT DARMSTADT

- New $c\bar{c}$ states at B factories: X, Y, Z
  (cf. Godfrey, arXiv:0910.3409)
  - Challenge for understanding of QCD
  - Unitary limit relevant?
- $X(3872)$ (Belle, CDF, BaBar, D0, LHCb)
- Nature of $X(3872)$?
  - $\bar{D}^0 D^{0*}$-molecule, tetraquark, charmonium hybrid, ...



$m_X = (3871.65 \pm 0.06)$ MeV, $\quad \Gamma = (1.19 \pm 0.21)$ MeV, $\quad J^{PC} = 1^{++}$ (PDG 2023)

- Assumption: $X(3872)$ is weakly-bound D^0-$\bar{D}^{0*}$-molecule

  $\Rightarrow \quad |X\rangle = (|D^0\bar{D}^{0*}\rangle + |\bar{D}^0 D^{0*}\rangle)/\sqrt{2}\,, \qquad B_X = (0.07 \pm 0.12)\ \text{MeV} \approx 1/(2\mu_{DD^*}a^2)$

  $\Rightarrow$ universal properties (Braaten et al., 2003-2008; ...)

Neutral charm mesons and $X(3872)$

TECHNISCHE UNIVERSITÄT DARMSTADT

- Approximate unparticles of three D^0/D^{0*} mesons
- Interaction of $X(3872)$ with D^0, $\bar{D}^0$, D^{0*}, $\bar{D}^{0*}$ determined by large a

(Canham, HWH, Springer, PRD **80**, 014009 (2009))

$$a_{D^0X} = -9.7a \qquad a_{D^{*0}X} = -16.6\,a$$

- Richer structure because of $X(3872)$ (bound state)

two charm mesons

three charm mesons



Neutral charm mesons and $X(3872)$

TECHNISCHE UNIVERSITÄT DARMSTADT

- Approximate unparticles of three D^0/D^{0*} mesons
- Interaction of $X(3872)$ with D^0, $\bar{D}^0$, D^{0*}, $\bar{D}^{0*}$ determined by large a

(Canham, HWH, Springer, PRD **80**, 014009 (2009))

$$a_{D^0X} = -9.7a \qquad a_{D^{*0}X} = -16.6\,a$$

- Richer structure because of $X(3872)$ (bound state)

two charm mesons

three charm mesons



Scaling Behavior

- Universal scaling for unparticles of three neutral charm mesons
  (Braaten, HWH, Phys. Rev. Lett. **128**, 032002 (2022))



XD point production



$$\frac{dR}{dE} \sim \left(E^{-(\Delta_1+\Delta_2-\Delta_3)/2}\right)^2 \sqrt{E} \approx E^{0.1} \qquad \text{for } E/|\varepsilon_X| \gg 1 \qquad \text{from conformal 3-pt. function}$$

$\Delta_1 = 3/2, \quad \Delta_2 = 2, \quad \Delta_3 \approx 3.10119/3.08697$ (Braaten, HWH, Phys. Rev. D **107**, 034017 (2023))

Scaling Behavior

- Universal scaling for unparticles of three neutral charm mesons
  (Braaten, HWH, Phys. Rev. Lett. **128**, 032002 (2022))



XD point production



XD elastic scattering

$$\frac{dR}{dE} \sim \left(E^{-(\Delta_1+\Delta_2-\Delta_3)/2}\right)^2\sqrt{E} \approx E^{0.1}, \qquad \sigma \sim E^{-1.6} \qquad \text{for } E/|\varepsilon_X| \gg 1$$

Δ_1=3/2, Δ_2=2, Δ_3≈3.10119/3.08697